

## 2

Draft : Environmental Impact Statement

**Chapter B7 Marine Ecology**

<b>B7.1</b>	<b>Relevance of Project to Marine Ecology</b>	<b>2</b>
<b>B7.2</b>	<b>Applicable Legislation and Policy</b>	<b>2</b>
<b>B7.3</b>	<b>Methodology</b>	<b>3</b>
B7.3.1	Nomenclature and Terminology	3
B7.3.2	Assessment Approach	5
<b>B7.4</b>	<b>Existing Environment</b>	<b>6</b>
B7.4.1	Background	6
B7.4.2	Marine Protected Areas and Wetlands of Conservation Significance	7
B7.4.3	Mangroves and Saltmarsh	9
B7.4.4	Seagrass Meadows	12
B7.4.5	Hard Benthic Substrata and Associated Communities	23
B7.4.6	Soft Sediment Habitat and Associated Communities	38
B7.4.7	Marine Species of Conservation Significance	52
B7.4.8	Exotic Marine Species and Pests	61
B7.4.9	Fish and Shellfish Resources	61
<b>B7.5</b>	<b>Assessment of Potential Impacts</b>	<b>73</b>
B7.5.1	Assessment Basis	72
B7.5.2	Overview	77
B7.5.3	Direct Modification of Benthic Habitats and Communities from Dredging, Dredged Material Placement and Wharf Upgrades	82
B7.5.4	Increased Suspended Sediment Concentrations and Sedimentation from Dredging and Dredged Material Placement	88
B7.5.5	Interactions between Marine Fauna and Vessels or Dredge Plant	108
B7.5.6	Acoustic Effects to Marine Fauna from Wharf Upgrade Works	110
B7.5.7	Other Indirect Interactions between Vessels and Marine Receptors	111
B7.5.8	Exposure of Marine Flora and Fauna to Debris, Spills and Dredging-Borne Contaminants	112
B7.5.9	Summary of Impacts to Marine Megafauna	114
<b>B7.6</b>	<b>Assessment Summary</b>	<b>115</b>
<b>B7.7</b>	<b>References</b>	<b>122</b>

## B7.1 Relevance of Project to Marine Ecology

This section describes the existing ecology of the marine environment in the vicinity of the Cairns Shipping Development Project (the project), and identifies potential impacts associated with the project. This chapter principally addresses Section 5.4.3 of the Environmental Impact Statement (EIS) Terms of Reference (TOR) (The Coordinator-General 2012); and Sections 5.6, 5.9 and relevant parts of Section 5.10 of the Commonwealth Guidelines for the Cairns Shipping Development Project EIS (DSEWPAC/GBRMPA 2013). This report section specifically describes:

- The main features of the existing environment in the study area, focusing on important or sensitive marine ecological resources and the integrity of coastal ecosystems
- Possible impacts on the ecology from dredging and placement of dredged material associated with the expansion of the shipping channel and swing basins
- Potential impacts on the marine environment from the construction and operation of the port facilities
- Options for managing and mitigating identified impacts.

In terms of impact mitigation, the project design has sought to avoid and/or minimise impacts to the greatest practicable extent. The mitigation strategies outlined in this chapter specifically aim to mitigate construction and operational impacts associated with the project.

## B7.2 Applicable Legislation and Policy

The following is a summary of Federal and Queensland legislation relevant to marine ecological aspects of the project.

### Federal:

- **Environment Protection and Biodiversity Conservation Act 1999** (EPBC Act), which provides for the protection of matters of national environmental significance (MNES). MNES of relevance to the project include:
  - World Heritage Properties – specifically Great Barrier Reef World Heritage Area (GBRWHA)
  - National Heritage Places – which encompass the entire area of the GBRWHA
  - Great Barrier Reef Marine Park (GBRMP) – adjacent to Cairns harbour; the GBRMP is managed under the Commonwealth Great Barrier Reef Act 1975
  - Commonwealth Marine Areas – which includes areas offshore from Cairns harbour
  - Nationally threatened species and ecological communities (including marine turtles and whales)
  - Migratory species (including dugong, whale shark, and several threatened marine mammal and turtle species)
  - The closest wetlands of international significance are Bowling Green Bay Ramsar site and Coral Sea Reserves Ramsar site; both located over 250 km from the Cairns port area. These are well outside the project's area of influence and are not considered further
  - **Great Barrier Reef Marine Park Act 1975** (GBRMP Act) – relevant to the protection of marine ecological values within the boundaries of the GBRMP that may be affected by the project.

### State:

- **Nature Conservation Act 1992** (NC Act), which provides for the protection of state-listed threatened and near-threatened species, which in the context of this project includes marine turtles, whales, dolphins and dugong
- **Marine Parks Act 2004**, which is relevant to the protection of marine ecological values within the boundaries of the state-controlled GBRMP
- **Fisheries Act 1994**, which provides for the use, conservation and enhancement of the fisheries resources and fish habitats. Of particular relevance to the project is the management of Fish Habitat Areas (FHA); the protection of fisheries habitats such as seagrass, mangroves and saltmarsh; and the protection of fish stocks
- **Environment Protection Act 1994** (EP Act), which provides for sustainable resource development while protecting ecological processes. The EP Act regulates environmentally relevant activities, including petroleum activities. The **Environmental Protection (Water) Policy 2009** aims to achieve the objectives of the EP Act in Queensland waters by establishing environmental values and water quality objectives
- **Queensland Coastal Plan** (Coastal Plan), which describes how the coastal zone is to be managed and identifies areas of high ecological significance (HES) and general ecological significance (GES), which includes wetlands adjacent to the project.

## B7.3 Methodology


### B7.3.1 Nomenclature and Terminology

For the purpose of this report the following terminology has been adopted (**Figure B7.3.1a**):

- The term *study area* refers to all waters within Cairns harbour, Trinity Inlet, and adjacent waters extending just past Double Island to the north and Cape Grafton to the east
- The *project areas* include:
  - *Terminal site* – refers to the Cairns Cruise Liner Terminal (CCLT) and the proposed minor wharf infrastructure upgrade construction footprint and immediate surrounds for the project, including existing cruise shipping wharves 1-5 and the construction footprint for associated ship services, including fuel supply, potable water and fire-fighting services
  - *Dredged material placement area* (DMPA) – the new offshore placement site proposed for this project
  - *Dredge footprint* refers to the outer channel, swing basins, inner channels and berth pockets that will be subject to capital dredging, and subsequent future maintenance, as a result of this project
- Existing offshore dredged material placement area refers to the existing offshore placement site.

The *surrounding area* refers to the intertidal and subtidal waters within 20 km of the study area.

Figure B7.3.1a Location of the Study Area and Project Areas



## B7.3.2 Assessment Approach

### B7.3.2.1 Existing Environment Methodology

Flora and fauna species, communities and habitats within the study area and surrounds were defined through searches of relevant databases, a review of previous studies, and where there was inadequate existing information, through supplementary field investigations. Searches were undertaken through the following databases: EPBC Protected Matters Search Tool; DEHP Wildlife Online database and DEHP StrandNet database. Relevant studies are cited in the text.

In order to source additional locally-specific information on marine megafauna in the Cairns area (e.g. turtles, dolphins, dugongs, etc.), consultation was undertaken with representatives from a range of state and federal government departments and local business operators, including:

- Mark Read (Great Barrier Reef Marine Park Authority).
- Ian Osbourne and Julie Dutoit (Department of National Parks, Recreation, Sport and Racing).
- Col Limpus and Justin Meager (Department of Environment and Heritage Protection).
- Jeffrey Krause (Department of Agriculture, Fisheries and Forestry).
- Noleen Brown (Department of Science, Information Technology, Innovation and the Arts).
- Dale Mundraby (Djunbunji Land and Sea Program).
- Isabel Beasley (James Cook University).
- Jennie Gilbert (Cairns Turtle Rehab Centre).
- Emma Scott (Wavelength Charters).
- Tim North (Reef Magic Cruises).

Information gaps were identified in the initial stages of the baseline reporting process; hence supplementary field surveys were undertaken to fill key gaps. The following field surveys were undertaken:

- Soft sediment habitat types and epifauna communities. Soft sediment habitats within and adjacent to the three project areas were mapped using acoustic techniques. Single beam sonar data were analysed using Qvester Tangent Corporation (QTC) software packages and a preliminary acoustic habitat class map was derived. Ground-truthing was then carried out to assess the sediment types within each acoustic habitat class, and to characterise epibenthic communities within each habitat class. Sediment types were sampled through grab based sampling and subsequently subject to sieve analysis to determine particle grain size distribution. Epibenthic communities were sampled using either a towed underwater video camera, or beam trawl, which was deployed at a total of 25 sites. The relative abundance of taxa recorded at each site (as recorded on a single transect line, standardised for four minutes) was then assessed
- Benthic infauna communities. A total of 24 sites were sampled at representative areas within the project areas and additional 'control' areas outside the disturbance footprint but within the study area and surrounds. Replicate 0.028 m<sup>2</sup> van Veen grab samples were collected at each site, sieved through a 0.5 mm screen and preserved in a 10 percent buffered formalin solution. Samples were sent to Stephen Cook (Dardanus Scientific) for identification and enumeration of each taxa. Summary statistics for abundance and taxa richness were derived, and multivariate statistical analysis was used to explore spatial patterns in community structure
- Shoreline Assessment. Intertidal communities were surveyed during spring low-tide periods to assess the major community constituents of rocky shores at East Trinity, False Cape, and Rocky Island
- Coral Surveys. Surveys of coral community composition and gross reef geomorphology were undertaken at Double Island and Rocky Island. Methods focused on i) quantifying benthic communities at various distances from the reef edge, and ii) mapping an indicative boundary of the reef edge and different reef habitats (i.e. ecotones). This was undertaken using a combination of photographic transects for Coral Point Count analysis, together with GPS and GIS mapping tools
- Baseline mapping of seagrass and benthic macroinvertebrate communities. James Cook University (JCU) conducted baseline mapping assessments of seagrass and macroinvertebrate communities specifically for this EIS. These assessments focused on Trinity Inlet, Cairns harbour and Trinity Bay. Specific methodologies can be found in the relevant JCU reports (McKenna et al. 2013, Rasheed et al. 2013, Jarvis et al. 2014).

A description of sampling and analysis methods, including maps of sampling locations, is provided in **Appendix D13**.

Extensive detailed seagrass and macrobenthic surveys were also undertaken by JCU's TropWATER throughout the Trinity Inlet and Cairns harbour area. Key findings of these surveys have been incorporated into the following existing environment description. The detailed TropWATER reports – including full methodology and results – are attached as **Appendix D13**.

### B7.3.2.2 Impact Assessment Methodology

The assessment of marine ecological impacts is based on a risk-based approach, taking into account consideration of the i) significance of a particular impact (i.e. intensity, scale and duration); ii) likelihood of that particular impact occurring; and iii) overall impact risk rating. This process is detailed in **Section B7.5**.

For potential marine ecology impacts relating to processes associated with water quality and sediment deposition, the results of the water quality modelling (**Chapter B5, Marine Water Quality**) and sediment deposition modelling (**Chapter B3, Coastal Processes** and **Appendix D4, Water Quality Model Development and Calibration Report**) undertaken specifically for this EIS have been used to predict marine ecology impacts. Specifically, this modelling has been used to predict the magnitude and location of particular water quality and sediment deposition impacts.

The modelling results, together with the ecological thresholds applied for this project, were used to infer the predicted location of ecologically meaningful 'zones of impact' (i.e. zone of high impact, zone of low to moderate impact, zone of influence). To determine the threshold values to delineate the zones of impact, a combination of water quality (turbidity) and biological tolerance methods was used. This entailed using baseline water quality monitoring data to set initial threshold values for impact zones. These threshold values and impact zone predictions were then tested using biological tolerances from literature values to check that the threshold values are biologically meaningful. This approach is described further in **Chapter B5, Marine Water Quality**.

## B7.4 Existing Environment

### B7.4.1 Background

Cairns harbour is a shallow, north-facing coastal embayment located adjacent to the city of Cairns. Cairns harbour is a low energy environment that is the receiving environment for several coastal drainages, including Trinity Inlet, Barron River and several coastal creeks. Due to its shallow depths and the plentiful supply of fine terrigenous sediments from the Barron River and other coastal drainages, wind and tidal currents regularly result in re-suspension of sediments and high turbidity within Trinity Bay.

Trinity Inlet is a large estuary that is fed by several minor drainages including Skeleton, Chinaman, Blackfellows, Wrights, Redbank, Wahday, Falls and Seelee creeks. It is thought that the Trinity Inlet once formed the mouth of the Mulgrave River, but was diverted southwards as a result of sediment accumulation on the coastal plain. As Trinity Inlet is not flushed by a major river, it represents a tidally dominant system with less variation in salinity than is usual in estuaries in high rainfall areas (Perry 1995). The coastal drainages, together with Admiralty Island, form a complex bio-physical setting that provides important habitats for a range of estuarine plant and animal species (Perry 1995; WBM 1997).

Cairns harbour and Trinity Inlet contain the following important ecological values:

- A wide diversity of marine habitat types including sandy beaches, mangrove forests, saltmarshes, intertidal shoals, seagrass meadows, subtidal soft sediment habitats, rock walls and rocky shores
- An extensive area of mangroves exhibiting a range of species and community types, some of which are limited in their distribution elsewhere
- Seagrass beds which represent one of the only two major seagrass areas between Hinchinbrook Island and Cooktown (however, the meadow size is presently far smaller than previously mapped - see **Section B7.4.4**)
- Mangroves, saltmarsh seagrass meadows and 'unvegetated' soft sediment habitats and other associated wetlands which have been recognised as important nursery areas for juvenile fish and prawns of commercial importance
- Habitats for a wide range of fish and shellfish species of direct economic significance
  - A range of habitat types that significantly underpin the biodiversity values of the region
  - Habitats that are important to migratory waders of international significance (see **Chapter B8, Terrestrial Ecology**)
  - Potential feeding areas for marine turtles, dugongs, whales, and dolphins, which are listed as threatened or migratory under Commonwealth and/or Queensland legislation.

This report section describes these environmental values and the condition/integrity of underpinning marine habitats (including benthic primary producer habitats such as seagrass, mangroves and saltmarsh communities) within the study area.


### **B7.4.2 Marine Protected Areas and Wetlands of Conservation Significance**

Protected and other ecologically significant areas represented within the study area include the following, as shown in **Figure B7.4.2a**:

- Areas designated as matters of national environmental significance under the EPBC Act, including the GBRMP) and GBR World Heritage Area (GBRWHA)
- Areas specifically protected under state legislation, namely Fish Habitat Areas (Trinity Inlet FHAs and other smaller FHAs located within the study area), GBR Coast Marine Park, GBR Wetlands, and wetlands listed as MSES (Cairns/ Trinity Inlet wetlands listed as a HES Wetland under MSES)
- Wetlands listed in the Directory of Important Wetlands in Australia (DIWA), which are not specifically protected under legislation but have recognised high conservation significance. Two DIWA wetlands occur in the study area: GBRMP and Port of Cairns/Trinity Inlet wetlands.

Information detailing the relevance of these protected areas and relevant management considerations are provided in **Chapter B2, Nature Conservation Areas**. Note that no Dugong Protection Areas occur within the study area, with the nearest being located approximately 150 km south at Hinchinbrook.

Figure B7.4.2a Location of Conservation Areas of Relevance to the Marine Ecology Aspects of the Project





## B7.4.3 Mangroves and Saltmarsh

### B7.4.3.1 General

At least 24 mangrove species from 15 genera have been recorded growing within Trinity Inlet as listed in **Table B7.4.3.1a** (Dowling 1983). No similar comprehensive inventory of saltmarsh species has been undertaken in the study area to date. Dominant saltmarsh species in the study area include *Sporobolus virginicus*, *Suaeda spp.*, *Sarcocornia quinqueflora* and *Tecticornia spp.* None of the mangrove or saltmarsh species occurring in the study area and surrounds are considered threatened or near threatened under legislation, and tropical saltmarsh and mangroves are not listed as threatened ecological communities under the EPBC Act.

Mangroves and saltmarshes represent benthic primary producer habitats, and support a range of functional ecosystem values. Mangrove and saltmarsh communities have a high ecological value as they provide food and shelter resources for a range of invertebrates, birds and fish (Robertson and Duke 1997). Many of the fish and shellfish species inhabiting mangals (mangrove forests) and saltmarshes are of direct recreational and commercial fisheries value. Mangroves and saltmarshes are also highly productive (Hutchings and Saenger 1987), are important in the stabilisation of the beds and banks of estuaries (Carlton 1974), and more recently, are increasingly being recognised for their high carbon sequestration potential. In accordance with these potential values, mangroves, saltmarshes and other marine plants are protected plants under the *Fisheries Act 1994* and a permit is required for their disturbance/removal.

**Table B7.4.3.1a Mangrove Species of Trinity Inlet (Adapted from Dowling 1983)**

Common Name	Species
holly-leaf mangrove	<i>Acanthus ilicifolius</i>
mangrove fern	<i>Acrostichum speciosum</i>
club mangrove	<i>Aegialitis annulata</i>
river mangrove	<i>Aegiceras corniculatum</i>
grey mangrove	<i>Avicennia marina</i>
ribbed-fruited orange mangrove	<i>Bruguiera exaristata</i>
large-leafed orange mangrove	<i>Bruguiera gymnorhiza</i>
small-leafed orange mangrove	<i>Bruguiera parviflora</i>
flat-leafed yellow mangrove	<i>Ceriops decandra</i>
yellow mangrove	<i>Ceriops australis</i>
long-fruited yellow mangrove	<i>Ceriops tagal</i>
wrinkle-pod mangrove	<i>Cynometra iripa</i>
milky mangrove	<i>Excoecaria agallocha</i>
looking-glass mangrove	<i>Heritiera littoralis</i>
red-flowered black mangrove	<i>Lumnitzera littorea</i>
white-flowered black mangrove	<i>Lumnitzera racemosa</i>
myrtle mangrove	<i>Osbornia octodonta</i>
tall-stilted red mangrove	<i>Rhizophora apiculata</i>
long-fruited red mangrove	<i>Rhizophora mucronata</i>
red mangrove	<i>Rhizophora stylosa</i>
yamstick mangrove	<i>Scyphiphora hydrophyllacea</i>
mangrove apple	<i>Sonneratia alba</i>
cedar mangrove	<i>Xylocarpus moluccensis</i>
cannonball mangrove	<i>Xylocarpus granatum</i>

### B7.4.3.2 Spatial Patterns

In Queensland's regional ecosystem (RE) mapping, mangrove forest is mapped as RE 7.1.1 (mangrove closed forest to open shrubland on areas subject to regular tidal inundation) and saltmarsh communities are comprised of RE 7.1.2 (*Sporobolus virginicus* grassland, samphire open forland and bare salt pans, on plains adjacent to mangroves). The distribution of these REs within the study area and surrounds is shown in **Figure B7.4.3.2a**, noting extensive mangrove forest occurring within Trinity Inlet and the less urbanised sections of Cairns harbour. These REs are listed as least-concern remnant vegetation under the *Vegetation Management Act 1999* (VM Act). Other coastal REs in the study area include dune and foreshore vegetation dominated by melaleucas, acacias and eucalypts (e.g. REs 7.2.3, 7.2.7, 7.2.8). These latter REs are typically of concern, with 7.2.7 and 7.2.8 listed as endangered (note that 7.2.7 fringes the coastline at Ellie Point). Refer to **Chapter 8, Terrestrial Ecology** for a description of RE types outside the intertidal zone.


There are three main estuarine wetlands within and directly adjacent to the study area that support remnant and regrowth mangrove forest and saltmarsh: Trinity Inlet/Admiralty Island, East Trinity and the Barron River estuary (**Figure B7.4.3.2a**). Saltmarsh communities are poorly developed and patchily distributed in the wet tropics as high rainfall limits the development of hyper-saline soils, allowing mangroves to dominate the intertidal zone (Smith and Duke 1987). Based on the RE mapping, the total area of mangrove forest in the study area (RE 7.1.1) was 4,420 hectares, whereas saltmarsh and saltpan covered an area of 257 hectares. There are no mangrove and saltmarsh areas within 100 m of the CCLT, dredging area or dredged material placement area (DMPA).

The distribution and community structure of mangroves and saltmarsh plants is controlled by four primary environmental factors: salinity, tidal range, degree of wave and current action, and the physical nature of the substrate (King 1981; Odum et al. 1985). Saltmarsh and mangroves grow in the intertidal zone, typically within calm, protected environments. The estuarine wetlands of the study area were comprised of a mosaic of community types, with the distribution and extent of communities varying among tidal zones. The following broad zonation patterns occur:

- Areas of closed *Aegiceras* (and in places) *Avicennia* +/- *Rhizophora* forest often formed a narrow band near the low tide mark
- Closed *Rhizophora* forest tended to numerically dominate areas towards the low water mark (typically landward of the *Aegiceras* and *Avicennia*), whereas mid-tide levels were dominated by a closed *Cerriops* forest
- Saltpan/saltmarsh formed small isolated patches in places. Saltmarsh areas were dominated by salt couch grasses (e.g. *Sporobolus virginicus*) and samphire (*Sarcocornia* spp).

Trinity Inlet has one of the most extensive coastal mangrove forests in the region between Port Hinchinbrook and Cooktown (WBM 1998). Elsewhere in the region, the steep terrain limits the development of extensive mangrove forests on the open coast (WBM 1998). However, estuaries in the region can contain mangrove forests that are similar in size and structure to those found in the study area (DEHP 2013).

**Figure B7.4.3.2a Remnant Mangrove and Saltmarsh Communities Based on Regional Ecosystem Mapping (Data source: DEHP 2013)**



### B7.4.3.3 Temporal Changes

The mangrove forests and saltmarshes of the study area have undergone extensive changes over time in response to land clearing/draining activities, and long-term changes in physical processes. Mangrove and claypan/saltmarsh habitats have been progressively removed in the broader Cairns harbour and Trinity Inlet area over the past few decades in association with airport development, agriculture (especially sugarcane farming on the eastern side of Trinity Inlet) and similar activities as the city of Cairns has developed. In contrast, a slight rise in sea level, together with increased sedimentation over the past 40 to 50 years (Goudkamp and Chin 2006), has led to increases in mangrove area within other parts of the study area. A comparison of modelled pre-clearing and recent (1999) regional ecosystem mapping for the Innisfail lowland subregion (including Cairns harbour and Trinity Inlet) indicates that the total area of mangroves has declined from 21,777 hectares to 9,078 hectares in 1999 (a net loss of 58 percent) (Accad and Neil 2006).

Similarly, claypan/saltmarsh communities were estimated to cover an area of 340 ha prior to European settlement, but now cover 65 ha (WBM 1997). This suggests that 80 percent of the original area has been lost over time. While clearing and draining were key contributors to saltmarsh/claypan losses, a landward expansion of mangroves has also resulted in net losses in saltmarsh/claypan extent. Olsen (1983) estimated approximately 40 percent of the saltmarsh/claypan area in Trinity Inlet was lost as a result of the landward progression of mangroves between 1952 and 1978.

### B7.4.3.4 Resistance, Resilience and Condition

Clearing and draining of freshwater wetlands, saltmarshes and mangrove forests has reduced the extent of these communities, and the loss of connectivity among terrestrial, freshwater and marine habitats (Perry 1995). Despite suffering major losses over time, the mangroves habitats of the study area are thought to be in good condition. Furthermore, rehabilitation and remediation works at the East Trinity Reserve have allowed mangroves and saltmarsh to recolonise areas previously severely affected by hydraulic modifications, clearing and exposure of acid sulfate soils (ASS) (refer regrowth values for East Trinity, **Table B7.4.3.4a**). New mangrove colonisation has also recently been evident progressing seaward adjacent the channel south of Bessie Point, extending south towards Stafford Point. The wetlands, while in a state of recovery, are providing locally important habitat for species of fisheries significance, as well as wader birds (see **Chapter B8, Terrestrial Ecology**).

Mangroves and saltmarsh are relatively tolerant of low to moderate increases in sediment loads, which is a necessary adaptation for living in a depositional environment. However, major increases in sediment loading can lead to burial of mangrove pneumatophores and saltmarsh plants, changes to sediment types, and/or changes to benthic fauna species, causing physiological stress and possible mortality to plants.

**Table B7.4.3.4a Area of Remnant Mangrove and Saltmarsh Regional Ecosystems (RE) Mapped in the Study Area (Data Source: DEHP 2013)**

Wetland	Mapped Area in Hectares of Remnant (and Regrowth) REs	
	Mangroves	Saltmarsh
Trinity Inlet/Admiralty Island	2,936 (<5)	216 (<1)
East Trinity	282 (270)	7 (83)
Barron River mouth	468 (<5)	18 (<1)

## B7.4.4 Seagrass Meadows

### B7.4.4.1 Spatial Patterns

A total of seven species of seagrass have been recorded in the study area (Coles et al. 1993; Rasheed et al. 2013), namely *Zostera muelleri* (formerly *Z. capricorni*), *Halodule uninervis*, *Halophila decipiens*, *Halophila ovalis*, *Cymodocea serrulata*, *Cymodocea rotundata* and *Thalassia hemprichii* (**Table B7.4.4.1a**). This compares with the 15 species known from the GBRWHA (UNESCO 2011).

Numerous studies have investigated the seagrass communities of Cairns harbour since approximately 1984 (Coles et al. 1995; Rasheed et al. 2013). Since 2001, the Queensland Department of Agriculture, Fisheries and Forestry (DAFF) or JCU - TropWATER have undertaken seagrass meadow surveys on an annual basis, which included sites located within the study area and surrounds. They represent the most comprehensive surveys undertaken in the area. The most recent JCU seagrass reports (Rasheed et al. 2013, Jarvis et al. 2014) are provided in **Appendix D13**, with key results mentioned below. For impact assessment purposes, the maximum historical extent of seagrass as shown in **Figure B7.4.4.1a** has been conservatively adopted as the baseline seagrass distribution.

Figure B7.4.4.1a Present (2013) Seagrass Distribution in Cairns Harbour and Trinity Inlet (Jarvis et al. 2014)




Table B7.4.4.1a Life-History Characteristics of Seagrass found in Cairns Harbour and Trinity Inlet

Species	Distribution <sup>2</sup>	Sensitivity to Disturbance	Reproductive Mode
<i>Zostera muelleri</i>	<ul style="list-style-type: none"> <li>• Among the largest <i>Zostera</i> beds in northern Queensland<sup>1</sup></li> <li>• Abundant in intertidal and shallow subtidal waters throughout Cairns Harbour.</li> </ul>	<ul style="list-style-type: none"> <li>• Large growing species providing large stores of primary energy reserves</li> <li>• High capacity to endure unfavourable conditions and tends to be stable in its distribution and abundance.</li> </ul>	<ul style="list-style-type: none"> <li>• Seed and vegetative</li> <li>• Moderate rhizome persistence, growth and reproductive output c.f. <i>Halophila</i> spp.</li> <li>• Recovery times longer than <i>Halophila</i> spp.</li> </ul>
<i>Halodule uninervis</i>	<ul style="list-style-type: none"> <li>• Generally co-dominant in intertidal and shallow subtidal waters with <i>Z. Muelleri</i></li> <li>• The dominant species in intertidal and subtidal areas on the eastern side of Cairns harbour (Bessie Point to False Cape).</li> </ul>	<ul style="list-style-type: none"> <li>• Small stores of energy reserves (owing to small plant size)</li> <li>• Rapidly declines when conditions become unfavourable for growth.</li> </ul>	<ul style="list-style-type: none"> <li>• Seed and vegetative</li> <li>• Fast growth and high reproductive output</li> </ul>
<i>Cymodocea serrulata</i>	<ul style="list-style-type: none"> <li>• Transient species, previously recorded in low abundance throughout Cairns harbour.</li> </ul>	As for <i>Z. muelleri</i> above.	<ul style="list-style-type: none"> <li>• Seed and vegetative</li> <li>• Low to moderate growth and reproductive output c.f. <i>Halophila</i> spp., high rhizome persistence</li> <li>• Recovery longer than <i>Halophila</i> spp.</li> </ul>
<i>Cymodocea rotundata</i>	<ul style="list-style-type: none"> <li>• Transient species, previously recorded in low densities near Ellie Point.</li> </ul>	As for <i>Z. muelleri</i> above.	As for <i>C. serrulata</i> above
<i>Thalassia hemprichii</i>	<ul style="list-style-type: none"> <li>• Recorded in Cairns harbour in 1993 but not thereafter.</li> </ul>	As for <i>Z. muelleri</i> above.	<ul style="list-style-type: none"> <li>• Seeds and vegetative</li> <li>• High rhizome persistence</li> <li>• Low growth</li> <li>• Seeds are buoyant and can disperse over long distances<sup>4</sup></li> </ul>
<i>Halophila ovalis</i>	<ul style="list-style-type: none"> <li>• Dominant in intertidal and shallow subtidal waters around Trinity Inlet and Cairns harbour</li> </ul>	<ul style="list-style-type: none"> <li>• Small stores of energy reserves (small plant size)</li> <li>• Rapidly declines when conditions become unfavourable for growth.</li> </ul>	<ul style="list-style-type: none"> <li>• Seeds and vegetative</li> <li>• Colonising growth strategy</li> <li>• Fast growth and high reproductive output</li> <li>• Seeds can remain viable in sediments for years<sup>3</sup></li> </ul>
<i>Halophila decipiens</i>	<ul style="list-style-type: none"> <li>• Subdominant in places along the mainland coast, and dominant (but sparse) in deep water environments.</li> </ul>	As for <i>H. ovalis</i> above.	<ul style="list-style-type: none"> <li>• Seeds</li> </ul> <p>As for <i>H. ovalis</i> for other characteristics</p>

1 = Coles et al. (1992); 2 = Rasheed et al. (2013); 3 = Rasheed and Taylor (2008); 4 = Olsen et al. 2004); 5 = Unsworth (2010)

Overall, up to a total area of 800 ha of seagrass habitat has been mapped in the study area and surrounds, with these meadows representing the only major seagrass resource between Hinchinbrook Island and Cooktown (Lee Long et al. 1993, 1996; Campbell et al. 2002, 2003; Jarvis et al. 2014). Historically, seagrass meadows have been recorded on shallow mud banks along the perimeter of Cairns harbour, and absent from deeper waters near the existing channel and the central section of Cairns harbour. Aggregated patches of seagrass have also been recorded in deeper waters north of False Cape and north-west of the existing channel, and along the foreshore of Trinity Inlet adjacent to Admiralty Island (mostly *Halophila* spp.).

Meadows tend to be denser and more structurally complex in the intertidal and shallow subtidal areas than those in deeper offshore waters (Rasheed et al. 2013). On the western side of the shipping channel, seagrass meadows have been dominated by *Zostera muelleri*, while *Halodule uninervis* has dominated the eastern side of the channel in the Bessie Point meadows. Narrow fringes of *Halophila* species have been found on the subtidal slopes of Trinity Inlet with *Zostera muelleri* found in some areas of the inlet on shallow intertidal/subtidal banks. Shallow waters favour the growth of larger growing species such as *Zostera muelleri* and *Halodule uninervis*, as well as *Halophila* species, which is a consistent pattern across most seagrass areas in North Queensland (Lee Long et al. 1993). The most well developed shallow water meadows were located along the foreshores on both sides of Cairns harbour (Rasheed et al. 2013).

*Halodule uninervis* has previously been detected as sparse isolated strands in the vicinity of the existing DMPA (Neil et al. 2003) and is infrequently observed within the channel and shallow marina berths by Ports North staff during annual monitoring of marine sediment and pests.


Seagrass meadows are not well established along the northern beaches or associated bays due to unsuitable habitat and substrate, however, a robust meadow is located around Double Island (refer Section B7.4.5.1).

**B7.4.4.2 Temporal Patterns**

Rasheed et al. (2013) and Jarvis et al (2014) collated monitoring data to assess spatial and temporal patterns in the structure of seagrass meadows within the study area. This data primarily comprised that from the long-term seagrass mapping conducted in the study area by the JCU researchers (formerly of DAFF), including mapping of all seagrasses in 1984 (Coles et al. 1985), 1988, 1993, 2001 and 2012; and annual monitoring of a subset of the seagrass meadows that has been carried out since 2001 (Figure B7.4.4.2b, Figure B7.4.4.2c). Monitoring of the percentage cover of seagrass at Ellie Point (Cairns harbour), carried out by Seagrass-Watch between 2001 and 2004, is also acknowledged. For the Seagrass-Watch monitoring, sampling was carried out on seven occasions and included both winter and summer sampling episodes; refer to Figure B7.4.4.2a for percentage cover estimates for representative sites (Johnson et al. 2011).

The results of these monitoring studies indicate the distribution, extent, biomass and density of seagrass assemblages within the study area can show great variation over a range of temporal scales. At seasonal scales, seagrass meadows in nearshore waters of the Great Barrier Reef region typically display a seasonal growth cycle in intertidal and shallow subtidal seagrass meadows (Waycott et al. 2005; Unsworth et al. 2009), with higher percentage cover of seagrass in late spring-summer than winter. McKenzie (1994) and Rasheed (1999) similarly describe distinct seasonality for seagrass communities at Cairns. Note that Seagrass-Watch (2013) monitoring data does not identify distinct seasonal patterns within the seagrass meadows of Cairns harbour, possibly reflecting the limited temporal replication of surveys (Figure B7.4.4.2a)

**Figure B7.4.4.2a Change in Seagrass Abundance (Percentage Cover) at Intertidal Meadows at Ellie Point from 2001 to 2004 (Source: Seagrass Watch 2013)**



Interacting with these general seasonal growth patterns are longer term (inter-annual) cyclic changes in seagrass meadows due to climate-driven disturbance and subsequent periods of recovery. Large inter-annual changes in seagrass meadow extent and community structure resulting from disturbance have been documented by Rasheed et al. (2013) and Jarvis et al. (2014). In summary, (**Figure B7.4.4.2d**, **Figure B7.4.4.2e**):

- Seagrass meadow extent showed an overall increase between 2001 and 2007, and a slight reduction in 2008 and 2009. This period was coincident with below average rainfall conditions
- There was a major reduction in seagrass above-ground biomass and extent between 2009 and 2010 sampling episodes. This was coincident with above average rainfall associated with 2010/11 La Niña events and Tropical Cyclone Yasi
- Surveys in 2011 and 2012 indicated there was a further decline in seagrass meadow extent and biomass. In November 2012 there was very little seagrass remaining within the study area. This is despite the study area experiencing favourable growing conditions in 2012
- Seagrass meadow extent in 2010-2012 was far below historical baseline levels observed during the 1980s and 1990s. These longer term (e.g. inter-annual, irregular) changes also interact with general seasonal growth patterns, resulting in a dynamic (and not necessarily predictable) community (**Figure B7.4.4.2f**).

The 2013 monitoring reported a very slight increase in both above ground biomass and total seagrass area since 2012. This presumably indicates the commencement of the recovery process.



Figure B7.4.4.2b Comparison of Seagrass Meadows in Cairns Harbour between Surveys in 1984, 1993, 2001 and 2012 (Source: Rasheed et al. 2013)




Figure B7.4.4.2c Comparison of Seagrass Meadows in Trinity Inlet between Surveys in 1984, 1993, 2001 and 2012 (Source: Rasheed et al. 2013)




Figure B7.4.4.2d Mean Above-ground Biomass (g DW m<sup>2</sup>) of all Monitoring Meadows Combined in Cairns Harbour and Trinity Inlet from 2001 – 2013 (Error Bars – Standard Error). Solid Blue Line Indicates 13-Year Mean of Meadow Biomass (Source: Jarvis et al. 2014)






Figure B7.4.4.2e Total Area of all Monitoring Meadows Combined in Cairns Harbour and Trinity Inlet from 2001 - 2013 (Error Bars – “R” Reliability Estimated). Solid Pink Line Indicates 13-Year Mean of Total Meadow Area (Source: Jarvis et al. 2014)



**Figure B7.4.4.2f Comparison of Seagrass Meadow Area in the Western and Eastern Sides of Cairns Harbour; 1984, 1988, 1993, 2001 and 2012 (Source: Rasheed et al. 2013)**



The combination of high rainfall, high river flow and cyclones can negatively impact seagrass either physically (burial, scouring, direct removal of plants and seed-banks) or physiologically (light limitation, excess nutrients and herbicides, and changes in salinity) (Rasheed et al. 2013). All years in the period 2007 to 2012 had above average rainfall, particularly in 2010 and 2011, while rainfall was below average in 2013 (Jarvis et al. 2014) (Refer **Appendix D13**). Barron River flow volumes were also well above average in 2007/08 and 2010/11. McKenzie et al. (2012) estimated that approximately 98 percent of intertidal seagrass areas in the regions directly affected by the path of Tropical Cyclone Yasi were lost as a consequence of destructive winds, and only a few isolated shoots remained in many coastal and reef habitats.

Light limitation is a key driver of spatial and temporal patterns in seagrass distribution and abundance, and is thought to be a key driver of the observed long-term temporal patterns in seagrass. As discussed in **Chapter 5, Marine Water Quality**, the light climate of the study area is a function of high total suspended sediment concentrations, plankton biomass and concentrations of dissolved organic carbon (e.g. tannins). Periods of high suspended sediment concentrations within the study area are controlled mainly by wave driven bed sediment remobilisation, and inputs of terrigenous sediments in flood waters from Barron River and the smaller coastal drainages entering Trinity Inlet and Cairns harbour (see **Chapter 5, Marine Water Quality** also **Appendix D13**). Conversely, when local climate conditions are in a drought-like state, subtidal seagrasses can thrive due to higher light levels reaching the seabed (Rasheed et al. 2013).

### B7.4.4.3 Seagrass Resistance, Resilience and Condition

Seagrass species differ in their sensitivity to disturbance (i.e. resistance) and capacity to recover following disturbance (i.e. resilience). The degree of resistance and resilience of seagrass depends on a number of often interactive factors including (Kenworthy 2000; Taylor and Rasheed 2009):

- Carbohydrate (energy) reserves to draw on during low light periods (resistance)
- Ability for photosystems to recover (resilience)
- Capacity for vegetative propagation (resilience)
- Seed bank occurrence (resilience)
- Historical and future disturbance regimes, including frequency, timing, duration and magnitude of disturbance (resistance and resilience).

**Table B7.4.4.1a** outlines the key life history characteristics of the seven seagrass species known from the study area. In general, small species such *Halophila* spp. tend to have low carbohydrate reserves compared to larger species, and are therefore the most sensitive species to low light conditions (i.e. low resilience). However, *Halophila* species also have adaptations that allow rapid recovery, including high reproductive output, rapid growth rates and the production of long-lived seeds that can remain viable in sediments for many years (i.e. high resilience). Larger species such as *Zostera muelleri* and *Cymodocea serrulata* have higher carbohydrate reserves and can therefore endure unfavourable periods for longer than small-bodied species (i.e. high resilience). However, these larger bodied species are slower to recover when lost (Rasheed 2004).


All seagrass species found in the study area are also capable of vegetative regrowth, which increases the capacity to recover following disturbance (i.e. rather than being solely dependent on sexual reproduction). Nonetheless, if all seagrass is lost from an area, some form of initial sexual colonisation would be required before seagrass is able to grow vegetatively. Recovery can be expected to be slower in instances where sexual colonisation is required to initiate the recovery process.

The capacity to recover following disturbance also depends on seagrass condition, which is a function of the previous disturbance history (magnitude, and spatial and temporal scale of disturbance). Successive periods of disturbance (i.e. multiple wet years) can deplete seagrass energy stores, seed banks and standing crop (i.e. seagrass condition), which greatly decreases the capacity for seagrasses to recover following disturbance. For example, the Reef Rescue Marine Monitoring Program (McKenzie et al. 2010) assessed the condition of seagrass meadows in the wider wet tropics region, with the sites nearest to the study area being located at Green Island. Overall, for the 2009/10 survey period, seagrass meadows of the wet tropics region were classified as being in a 'poor state' in coastal intertidal habitats, and a 'moderate state' at intertidal reef habitats (McKenzie et al. 2010).

Jarvis et al. (2014) applied a recently developed condition index to monitoring meadows, based on the results of annual monitoring of seagrass biomass, area and species composition. **Figure B7.4.4.3a** presents a summary of the present condition of seagrass monitoring meadows in the harbour and Trinity Inlet. All meadows in the harbour were rated as being in 'poor' condition on account of poor biomass and area; while meadows in Trinity Inlet ranged from 'poor' to 'good' condition.

In summary, seagrasses in the Cairns harbour are currently in a poor state and highly vulnerable to further declines. The next few growing seasons will be critical for the recovery trajectory of seagrasses in Cairns since propagule sources for recolonisation are limited and seed banks are ageing with reduced seed viability. This means that it will be critical to maintain conditions suitable for seagrass growth over the coming years.

Figure B7.4.4.3a Seagrass Monitoring Meadow Condition in Cairns Harbour, October 2013 (Jarvis et al. 2014)



## B7.4.5 Hard Benthic Substrata and Associated Communities

### B7.4.5.1 Habitats and Communities

Coral reefs form a benthic primary producer habitat. Based on mapping from the GBRMPA Gazetteer (**Table B7.4.5.1a**; **Figure B7.4.5.1d**), the total area of reef habitat within the study area is approximately 196.3 hectares. It includes mainland shallow fringing reefs and rocky shores, and well-developed reef platforms around nearshore islands. Habitat characteristics of reefs vary greatly among locations, reflecting differences in reef morphology.

#### Mainland Reefs and Rocky Shores Cairns Harbour/Yorkeys Knob

Within the harbour and Trinity Inlet, rocky shores and boulders are present along sections of the coastline near East Trinity (Bessie Point, Lyons Point) and surrounding False Cape. Rocky shores of East Trinity are mostly located in the intertidal zone, and consist of granite boulders and small outcrops surrounded by sand and mud substrata. The sedimentary coastline along the western side of the harbour and inlet are not known to support rocky shores. Site inspections undertaken as part of this EIS indicated rocky shore communities here were comprised of species typically found in turbid nearshore environments. At East Trinity, the granite boulders of the upper intertidal zone were densely covered in oysters (**Figure B7.4.5.1c -A**) with occasional low, closed mangroves present in places. The lower intertidal rock substrata were often covered in a dense mat of calcareous tubeworms (**Figure B7.4.5.1c -B**) with occasional clusters of brown macroalgae (**Figure B7.4.5.1c -E**) and a fine turf of filamentous green algae. Filter-feeding holothurians were the most common large conspicuous invertebrates observed (**Figure B7.4.5.1c -C**) with occasional soft corals (**Figure B7.4.5.1c -D**) and large molluscs also present (**Figure B7.4.5.1c-F**).

The rocky shores surrounding False Cape provide a more substantial subtidal habitat than East Trinity, with continuous rocky shores along a steeper shoreline profile (**Figure B7.4.5.1c -G**). A variety of macroalgal species including *Halimeda* and *Sargassum* (**Figure B7.4.5.1c -J**) dominated the lower intertidal area, with much of it covered in a thick turf of algae, encrusting sponges and bryozoans. Occasional encrusting faviid hard corals (**Figure B7.4.5.1c -I**) were present but uncommon (<five percent cover).

**Table B7.4.5.1a Reefs Mapped Within the Study Area and Surrounding Coastal Areas by the Great Barrier Reef Gazetteer**

Locality	GBRMPA Gazetteer Name and Number	Area (ha)
<b>Study Area</b>		
Mission Bay	Rocky Island Reef 06-051	0.2
	U/N Reef 16-050	10.57
Cooks Bay South	U/N Reef 16-094	0.89
Double Island	Haycock Reef 16-048	88.47
	Double Island Reef 16-047	96.19
<b>Surrounding Areas</b>		
Palm Cove to Port Douglas	U/N Reef 16-093	2.96
	U/N Reef 16-092	25.68
	U/N Reef 16-091	2.45
	Unity Reef No. 1 16-045A	9.03
	Unity Reef No. 2 16-045B	8.54
	U/N Reef 16-090	2.27
	U/N Reef 16-089	17.8
	Garioch Reef 16-082	20.49
	Alexandra Reefs (No. 2) 16-039B	411.74
	Yule Reef 16-081	11.13
	Korea Reef 16-080	208.98
	Alexandra Reefs (No. 1) 16-039A	538.11
	Wentworth Reef 16-037	104.91
	Egmont Reef 16-038	10.68

Locality	GBRMPA Gazetteer Name and Number	Area (ha)
Cape Grafton	U/N Reef 16-095	0.06
	U/N Reef 16-096	0.33
	U/N Reef 16-097	0.09
	U/N Reef 16-098	1.21
	U/N Reef 16-099	0.89
	U/N Reef 16-100	1.38
	U/N Reef 16-102	0.26
	U/N Reef 16-103	0.86
	U/N Reef 16-104	0.08
	U/N Reef 16-105	0.47
	U/N Reef 16-106	1.17
Turtle Bay/Little Turtle Bay/Wide Bay	U/N Reef 16-107	0.2
	U/N Reef 16-108	2.59
	U/N Reef 16-109	0.61
	U/N Reef 16-110	2.29
	U/N Reef 16-052	7.89
	U/N Reef 16-053	14.46
	U/N Reef 16-111	6.76
	U/N Reef 16-112	21.14
	U/N Reef 16-113	2.29
	U/N Reef 16-056	6.04
	U/N Reef 16-801/Gunjarra Island	17.31
	U/N Reef 17-074	0.43
	U/N Reef 17-075	0.36
U/N Reef 17-076	0.88	
Fitzroy Island	Little Fitzroy Reef 16-055	10.47
	U/N Reef 17-054G	4.64
	U/N Reef 17-054F	17.32
	U/N Reef 17-054C	0.61
	U/N Reef 17-054E	0.3
	Fitzroy Island Reef (No. 2) 16-054B	2.49
	Fitzroy Island Reef (No. 1) 16-054A	16.39

Further afield, Yorkeys Knob located approximately nine km north of Cairns harbour is a rock promontory that contains steep-gradient, plunging rocky shores with limited erosional bench or platform. Several rock promontories separated by a series of beaches occur along the coast to the north of Yorkeys Knob (e.g. Taylor Point at Trinity Beach and Buchan Point at Palm Cove). Taylor Point, located approximately 12 km north of Cairns, contains a small (0.89 ha) fringing reef that is mapped in the GBR Gazetteer (Reef 16-094).



## Double Island

Double Island (Double Island and Haycock Reefs: GBR Gazetteer number 16-047 and 16-048) contains a narrow fringing reef to the north and an extensive reef platform to the south, and has a total mapped area of 185 ha. This represents the largest reef in the study area (**Table B7.4.5.1a**).

The reef flat was comprised of a broad inner sediment zone that was surrounded by a rim of corals and algae along the reef front. The inner sections of the reef flat contained a mosaic of seagrass, macroalgae and bare sediments. A large sand spit extended in a north-south orientation along the north-western part of the reef flat.

Seagrass communities tended to dominate sandy substrates across the reef flat, whereas rubble substrates on and surrounding the reef flat were dominated by a variety of hard and soft corals and macroalgae. Extensive seagrass meadows were present on the reef flat east and south of the sand spit, whereas to the west of the spit, seagrass meadows were sparse and patchy. Seagrass communities were comprised of *Halophila ovalis*, *Halodule uninervis* [thin and thick forms], *Cymodocea serratula*, *Syringodium isoitefolia*. Seagrass cover and community composition was variable across the flat, but reached up to 70 percent in places. Dense meadows were dominated by *C. serratula* and *H. uninervis*.

The reef flat (particularly in the south-west) had numerous low-profile microatolls, consisting of *Porites* (massive and digitate forms), *Acropora*, and *Goniastrea*, but had low coral cover overall. Macroalgae and bare substrate was also abundant.

Quantitative transects located on the upper slope were dominated by bare substrate and in places hard corals, whereas corals and macroalgae co-dominated the deeper sections of the reef slope. The densest coral cover occurred on the reef front, with an overall average hard coral cover of 22.7 percent  $\pm$  3.58 (S.E.) and average soft coral cover of 8.2 percent  $\pm$  2.60 (S.E.). Hard coral cover was highest at the south-eastern edge of the reef (DI1-2; 36 percent), while the south-western reef edge (DI9-10) had high soft coral cover (17 percent) but lower hard coral cover (12.5 percent). The other surveyed sections of the reef slope had intermediate levels of hard and soft coral cover.

The dominant coral species varied among and within sites (**Figure B7.4.5.1a**). Within locations, *Acropora* numerically dominated on the deep slope transect (three-17 percent cover), but with the exception of DI.7 on the western reef margin (19 percent), generally had low cover on upper slope transects (0-one percent). *Montipora*, *Porites* and *Mycedium* were locally abundant, again more so in the deep slope than shallow slope transects. The relative abundance of soft corals varied marked across the reef, with *Efflatounaria* abundant at the north-west edge of the reef (DI9-10), and *Sinularia* abundant on the western reef edge (DI7-8).

Coral taxa (genera) richness varied marked among and within sites (**Table B7.4.5.1b**). DI1-2 on the south-eastern edge had high hard (10 taxa) and soft (five taxa) coral richness. The northern fringing reef (DI5-6) and southern-western edge (DI9-10) had moderate to high hard coral richness (12 and seven genera, respectively), but only one=two soft coral genera.

Macroalgae were numerically dominant across the reef flat and along the ridge of reef edge (**Figure B7.4.5.1a**). Turfing algae was abundant on all transects, ranging from 13-52 percent cover (average = 26.6 percent  $\pm$  4.47 S.E.). The canopy forming *Sargassum* was locally abundant, particularly on the western margin (DI7-8) of the reef flat.

**Figure B7.4.5.1b** is a non-metric, multi-dimensional scaling ordination showing patterns in reef community similarity at Double Island Reef and Rocky Island. There was great spatial variability in communities at Double Island Reef, whereas those at Rocky Island were more uniform and distinctly different from those at Double Island Reef.

Overall, the reef community at Double Island was in excellent condition, particularly on the reef slope and areas mapped as 'extensive hard and soft coral, sparse sargassum' and 'digitate *Porites* and *Acropora* with macroalgae'. Across these areas, corals were diverse in terms of species richness and in excellent condition, particularly for an inshore reef so close to the mainland. Being located inshore, most well-established species present at Double Island would have some resilience to windows of increased turbidity, to the extent that turbidity is increased during natural disturbance events in the northern beaches area.

**Table B7.4.5.1b Numbers of Coral General Recorded at each Transect and Site**

Location	Site	Transect	Hard corals	Soft corals
Double Island	DI.A	DI.1	6	3
		DI.2	5	3
		Sum	10	5
	DI.B	DI.3	2	0
		DI.4	2	1
		Sum	3	1
	DI.C	DI.5	8	1
		DI.6	2	1
		Sum	12	2
	DI.D	DI.7	2	1
		DI.8	2	1
		Sum	3	1
	DI.E	DI.9	3	1
		DI.10	3	1
		Sum	7	1
Rocky Island	RKY.A	RKY.1	2	1
		RKY.2	1	1
		RKY.3	0	1
		Sum	2	1
	RKY.B	RKY.4	2	0
		RKY.5	4	1
		RKY.6	2	1
		Sum	5	1
	RKY.C	RKY.7	4	2
		RKY.8	1	2
		RKY.9	0	1
		Sum	4	2

**Figure B7.4.5.1a Percentage Cover of (i) Broad Benthic Groups (top) and (ii) Numerically Dominant Taxa (bottom).**  
**Note: Odd Number Transects at Double Island were Located on the Upper Edge of the Reef Slope, even Number Transects were Located on the Deep Margin of the Reef Slope**





Figure B7.4.5.1b Non-metric MDS Ordination – Reef Communities



### Rocky Island/Mission Bay

The only reefs and rocky shores mapped by the GBRMPA Gazetteer within the southern parts of the study area are Rocky Island Reef and an associated unnamed fringing reef in Mission Bay (Figure B7.4.5.1d).

Site inspections at intertidal platform reef surrounding Rocky Island identified a diverse assemblage of corals, algae, large vertebrates and invertebrates within a relatively small area on the north-west extremity of the reef. The reef was composed of granite boulders surrounding most of the island, covered by barnacles and oysters in the upper intertidal zone (Figure B7.4.5.1e-A).

Coral cover was highest along the reef edge and attenuated to very low cover within approximately 50 m of the reef edge. Beyond the reef edge, the inter-reefal substrate consists of muddy sand interspersed with micro-atolls (Figure B7.4.5.1e-C). Hard coral genera including *Turbinaria* (Figure B7.4.5.1e-D), *Galaxea* (Figure B7.4.5.1e-B), *Favia* (Figure B7.4.5.1e-E), *Porites*, *Goniastrea*, and *Favites* dominant. The soft coral *Briareum* (Figure B7.4.5.1e-C) was very abundant, while soft corals *Cladiella* and *Sinularia* (Figure B7.4.5.1e-D) were found occasionally. Hard coral genera including *Dendrophyllids* (Figure B7.4.5.1e-F), *Platygyra*, *Montipora*, and *Moseleya* were present in low numbers.

Generally speaking, highest percentage of living coral was present along the southern and eastern faces of the reef. Water clarity was reasonably turbid along the northern reef edge, despite the extremely calm conditions and lack of significant rainfall. The sea floor was extremely silty and contained more of the species associated with depositional environments such as sea whips, (*Junceella* sp.) and soft corals. This part of the reef edge was most sheltered from south-east trades given its orientation and proximity to the lee of the continental part of the island. The western and north-western reef edge was shallower than the southern and eastern reef slope and this was probably due to accretion from terrigenous sediments, perhaps due to remobilisation by refracted waves.

During the site assessment, numerous molluscs, crustaceans, and large vertebrates were sighted including epaulette sharks (Figure B7.4.5.1e-G), three green turtles resting on the reef (Figure B7.4.5.1e-H), and several ornate crayfish (Figure B7.4.5.1e-I). Central parts of the reef flat contained a large number of broken fragments of staghorn (*Acropora*) coral (Figure B7.4.5.1e-J).

In contrast to Double Island, the reef community at Rocky Island appeared to be in a degraded condition, as evidenced by the high extent of old dead corals present, compared with the smaller and sparse extent of live coral. Coral species remaining on Rocky Island tended to be those most resilient to environmental and physical disturbances (including, for example, windows of increased turbidity). However, this community may also be under significant stress, given its present condition.

### *Artificial Hard Substrate*

In addition to natural reef systems, Trinity Inlet contains large areas of rock wall habitat, particularly within the port area and along the foreshore of Cairns city. Site inspections undertaken in the present study suggest the existing rock walls around the port support algal and invertebrate (primarily sponge and hydrozoan-dominated) communities. Rock wall habitats represent aggregation areas and habitat for a range of fish and shellfish species, and as such, can represent locally important fisheries habitats. Anecdotal observations of large numbers of boat-based recreational anglers adjacent to rock walls and other structures suggest that this area supports locally important fisheries habitats.

### *Surrounding Areas*

The GBRMPA Gazetteer also identifies a series of unnamed rocky shores along the east side of Cape Grafton, several of which support fringing reefs. Fitzroy Island, located offshore from Cape Grafton, also supports well developed fringing reefs and rocky shores.

Mid-shelf reef systems (including Green Island, the Arlington Reef complex, etc.) are located more than 20 km to the north-east of the study area.

Figure B7.4.5.1c East Trinity (A-F) and False Cape (G-J) Shoreline Assessment: Stilt Mangroves (*Rhizophora stylosa*) and Dense Oysters in the Upper Intertidal Zone (A); Encrusting Tubeworms cover the Lower Intertidal Zone(B); Filter Feeding Holothurians (*Pseudocolchirus violaceus*) were Common (C); Dendronephthyid Soft Coral (D); Brown Macro Algae (E); the Baler Shell (*Melo amphora*) (F) ; Dense Boulder Field (G); Turfing Algae and *Halimeda* (H); Hard Coral (*Favia stelligera*) (I) and *Sargassum* (J).




Figure B7.4.5.1d GBRMP Gazetteer Reefs in the Surrounding Area




Figure B7.4.5.1e Reef Communities at Rocky Island: Looking South across the Reef Flat (A); *Halimeda*, the Hard Coral *Galaxea* (B) and Soft Coral (*Briareum*) (C) were Dominant along the Reef Edge; Microatoll with *Turbinaria* and *Sinularia* (D); Hard Coral (*Goniopora*) (E); Dendrophyllid Hard Coral (F); Epaulette Shark (*Hemiscyllium ocellatum*) (G); Green Turtle (*Chelonia mydas*) (H); Ornate Crayfish (*Panulirus ornatus*) (I); and Bare Staghorn Coral (*Acropora*) Remnant Substrate (J).





Figure B7.4.5.1f Coral, Algae and Seagrass Communities at Double Island Reef





Figure B7.4.5.1g Coral, Algae and Seagrass Communities at Rocky Island Reef



**Figure B7.4.5.1h Reef Communities at Double Island Reef: Sparse Seagrass over Sand (A); Dense Seagrass with Macroalgae and Occasional Coral (B); Macroalgae Dominant with Seagrass and Coral (C); Macroalgae Dominant (Particularly Sargassum), Rubble and Hard Coral (D); Branching Portitid and Acroporid Hard Corals with Rubble (E); Extensive Hard and Soft Coral Communities (F) on the reef slope.**





Figure B7.4.5.1i Sample Photos of Extensive Hard Coral Communities at the Double Island Reef Slope



**Figure B7.4.5.1j Reef Communities at Double Island Reef: Dense Macroalgal Cover Over Rock and Rubble (A); Sparse Seagrass over Sand (B); Macroalgae and Occasional Soft Coral over Sand (C); Rubble with Macroalgae and Occasional Hard Coral (D); Macroalgae Dominant with Seagrass and Coral (C); Macroalgae Dominant (Particularly *Sargassum*), Rubble and Hard Coral (D); Hard Coral Microatolls with Soft Coral, Sand and Algae (E, F)**



### B7.4.5.2 Key Drivers

The Cairns harbour receives an abundant supply of alluvial sediments, and has high turbidity and sedimentation levels (see **Chapter 5, Marine Water Quality**). Furthermore, it experiences periods of low salinity during floods (Coles et al. 1987). These are key drivers of the structure of rocky shore and reef communities, and in the case of sediment supply, the distribution and extent of hard substrate habitat. In this regard:

- No low-lying rocky reef or fringing coral reefs were present in the harbour. Exposed rock substrate was comprised entirely of massive boulders and rocky shore outcrops within the (hydro-dynamically active) intertidal zone
- Autotrophic species (i.e. macroalgae, hard corals) were poorly developed in the harbour and were primarily comprised of small, isolated colonies.

The few coral species observed on rocky shores in the harbour have adaptations that allow them to cope with high sediment loads. This includes for example, (i) the capacity for some corals to switch from phototrophic to heterotrophic feeding strategies by feeding on suspended sediments; (ii) rapid replenishment of energy reserves between turbidity events; (iii) rapid rates of photo-acclimation; and (iv) energy conservation through reduced respiratory and excretory losses (Anthony and Larcombe 2000). Many nearshore species also secrete mucus to eliminate settled sediment.

### B7.4.6 Soft Sediment Habitat and Associated Communities

#### B7.4.6.1 Distribution of Habitats

Soft sediment habitats within Trinity Inlet and Trinity Bay include sandy beaches, intertidal mudflats and subtidal soft sediments. Soft sediment habitats in shallow areas (where there is sufficient light to allow photosynthesis) contain microphytobenthos assemblages, which are often important drivers in coastal food chains. The extent of 'unvegetated' soft sediment habitats can show great temporal variability in response to temporal changes in the extent of seagrass meadows (see **Section B7.4.4**).


Surveys carried out to map soft sediment habitats surrounding the proposed terminal site, dredge area and existing and potential DMPAs showed the seafloor was comprised of numerous substrate classes (**Figure B7.4.6.1a**). These classes were for the most part, serially ordinated over a large geographic area, with class 1-4 substrata found in inshore embayments, graduating to higher classes with distance offshore and increasing depth. Fine-scale heterogeneity in sediment types was found inside Trinity Inlet, in hydrological diverse channel areas and at the existing DMPA, which has received dredged material from elsewhere. Apart from these areas, changes in sediment classes occurred gradually over large geographic areas.

Summary particle size data for the acoustically defined substrate classes are shown in **Figure B7.4.6.1b**. The distribution of silts and clay (fines) with respect to distance from shore followed a bell-shaped distribution, with the least amount of fines present in inshore (classes one-four) and offshore areas (classes 10-12), with the midshore region (classes six-nine) dominated by fines. While the inshore and offshore classes had similar particle size distributions, the nature of the sands and coarser material between these two areas differed greatly (see also **Figure B7.4.6.1c**). Class 1 sediments consisted of fine, muddy, well-sorted sands, often with regular bedforms. Offshore classes 10, 11 and 12 had sand fractions consisting of large amounts of shell and shell grit. Video observations of these offshore areas also showed occasional patches of rubble.


The colour of the offshore material was also much lighter than material in the mid-shore regions, which was probably related to high concentrations of calcium carbonate. Offshore samples were much closer to Green Island; hence, have greater exposure to calcium carbonate originating from coral reefs.

It should be noted that while the acoustic classification resolved 12 classes of sediment, the biological relevance of fine-scale changes in PSD between adjacent classes was in many cases very low (see below). In each sediment type, fines made up a significant proportion of the substrate, and a silt layer was ubiquitous over the sea floor. The acoustic substrate mapping was able to distinguish very fine-scale differences in sediment type, but in reality, each sediment class consisted of varying degrees of sand and mud, with occasional gravel. Soft sediments dominate vast parts of the study area, with reef, cobble and rock wall habitat contributing very little to the overall areas.

Figure B7.4.6.1a Interpolated Soft Sediment Habitat Classes



**Figure B7.4.6.1b Particle Size Distributions for a Selection of Acoustic Substrate Classes.**  
 Class Colours Correspond to Figure B7.4.5.1d



Overall, the sediment classes detected by the acoustic surveys, together with intertidal sediments, can be grouped into four broad sedimentary habitat types across the study area, including:

- Intertidal banks and mud flats – Primarily concentrated around the eastern and western sides of the Cairns harbour, Yorkeys Knob and the Barron River mouth, and southern Mission Bay
- Subtidal inshore fine sandy mud – Distributed across subtidal sections of the harbour, Mission Bay, and remaining inshore marine reaches along the south western extent of the study area
- Midshore mud – The dominant broad soft sediment habitat type within the study area, covering an area of 10,140ha (**Table B7.4.6.1a**), including most of the northern section of the study area
- Offshore muddy sand and shell – Confined to the north eastern corner of the study area, which extends into the deeper, offshore bathymetric contour.

**Figure B7.4.6.1d** provides an indicative summary of the distribution of these broad sedimentary habitat types, with total areas within the study area listed in **Table B7.4.6.1a** (note: excludes muds in Trinity Inlet beyond the acoustic survey extent). Beyond the study area, broad sedimentary habitat types were largely dominated by: the ‘mid-shore mud’, extending across both near-shore and mid-shore waters; and ‘offshore muddy sand and shell’ in deeper offshore areas. The existing DMPA, together with DMPA Options 1, 2 and 3, are all largely comprised of the broad mid-shore mud sediments.

**Table B7.4.6.1a Area of Broad Soft Sediment Habitat Types Mapped in the Study Area**

	Area (ha)
Intertidal flats	2,142
Subtidal inshore fine sandy mud	6,131
Mid-shore mud	10,140
Offshore muddy sand and shell grit	1,267



Figure B7.4.6.1c Photos of Sediment from a Selection of Sediment Classes. Class Colours Correspond to Figure B7.4.6.1b, and Site Tags have been removed to allay confusion between Site Numbers and Sediment Classes.





Figure B7.4.6.1d Distribution of Major Soft Sediment Typ




### B7.4.6.2 Epibenthic Fauna


Video-based surveys undertaken simultaneously with the acoustic habitat mapping suggested epibenthic communities were sparse and patchy throughout the study area. Of the 14 useable video transects, seagrass was not observed and epifauna were recorded on 12 of the 14 transects. In total, 11 fauna taxa were recorded consisting of 64 animals, without seagrass or macroalgae.

Epibenthos assemblages observable in videos were dominated by burrowing gobies, sea pens (Figure B7.4.6.2a-A) and crinoids (feather stars, Figure B7.4.6.2a-B). Half of all specimens observed were burrowing gobies. There were 13 sea pens and seven crinoids. There tended to be more taxa (Figure B7.4.6.2b), more individuals (Figure B7.4.6.2b), and more burrows (Figure B7.4.6.2c) at sites located farther from shore (higher sediment classes) than in sites closer to shore.


**Figure B7.4.6.2a Small Sea Pens (A) and Feather Stars (B) were the most Abundant Sessile Taxa**



**Figure B7.4.6.2b Mean Abundances (per Four Minute Video Transect) of Epibenthic Fauna at Representative Sediment Classes**



**Figure B7.4.6.2c Mean Burrow Abundances (from three Frame-grabs per Transect) at Representative Sediment Classes**



McKenna et al. (2013) surveyed epibenthic macroinvertebrates, seagrass and macroalgae from 572 sites over the study area between October 2012 and January 2013 (**Figure B7.4.6.2d**; refer also full report in **Appendix D13**). Similar to findings of the acoustic habitat validation, McKenna et al. (2013) reported habitat forming macro-invertebrates such as soft corals and bryozoans were predominantly found further offshore, with little to no epibenthos found in the inner port or Trinity Inlet. Overall, McKenna et al. (2013) found the dominant habitat feature to be open mud/sand substrate with a low cover of epibenthic life from a diverse array of taxa. There were no high density benthic communities (>80 percent cover), and there were no unique or unusual benthic macroinvertebrate or algal communities found within the survey area. Epibenthic invertebrates were typically observed as isolated individuals, with the exception of:

- High density soft coral and seapens at two locations (one near west of the shipping channel and one in the vicinity of the DMPA – refer highest density polygons on Figure 7.4za).
- Medium density polychaetes at one area (north of the end of the shipping channel), echinoids at one area (south of the DMPA), and bivalves along the inshore area between Bessie Point and False Cape.

Epibenthos density polygons are shown overlaying the acoustically defined sediment class map in **Figure B7.4.6.2e**. A description of the various epibenthic macroinvertebrate regions identified by McKenna et al. (2013) is also provided in **Table B7.4.6.2b**.

Benthic macroalgae was shown to have a wide distribution, occurring in isolated areas throughout the harbour and Trinity Inlet (McKenna et al. 2013). Despite the wide distribution, percentage cover of algae was typically low, with 77 percent of macroalgae regions mapped having less than 10 percent cover. No high density macroalgae coverage was recorded. Macroalgae was generally dominated by *Sargassum* spp., *Halimeda* spp., as well as various mosaics of unidentified turfing algae mats and filamentous algae. Algae concentrations tended to occur in areas of the inner port and Trinity Inlet rather than offshore, particularly along the western harbour adjacent to the Esplanade. Comparisons with past data indicated that macroalgae distribution and density can fluctuate greatly (**Figure B7.4.6.2f**), partly in response to water quality (e.g. nutrients) and climatic factors (e.g. physical disturbance), as well as in response to changes in competitive phyto-organisms (i.e. opportunistic growth following reductions in seagrass extent or condition) (McKenna et al. 2013).

Many of the algae and benthic macroinvertebrate communities described in their survey occurred in proximity to maintained channels, port facilities and the DMPA.

Figure B7.4.6.2d Extent and Distribution of TropWATER Benthic Macroinvertebrate Survey Sites (from McKenna et al.2013)






Figure B7.4.6.2e Interpolated Sediment Classes with JCU Epibenthos Densities (McKenna et al. et al. 2013)



**Table B7.4.6.2b Epibenthic Macroinvertebrate Regions and Descriptions within Cairns Harbour and Trinity Inlet, refer Appendix D13 for Map of Region Locations (Source: McKenna et al. 2013)**

Density category	Benthic macro-invertebrate region description	Region ID (see maps 2 & 3)	No. of sites	Area $\pm$ R (ha)
Open substrate, occasional benthic individuals	Open substrate with isolated benthic individuals	1 (white survey sites on maps 2 & 3)	517	na
Low density benthic community	mostly open substrate with patches of low density bivalves & low numbers of individual polychaetes	2	2	6.59 $\pm$ 0.89
	mostly open substrate with patches of low density sea pens, hydroids and ascidians with low individual numbers of other taxa	3 & 6	9	1482.69 $\pm$ 561.99
	mostly open substrate with patches of low density bryozoans and hydroids and low individual numbers of other taxa	4 & 5	19	2248.81 $\pm$ 1730.27
	mostly open substrate with patches of low density sea pens, hydroids and ascidians with moderate numbers of individual polychaetes & low numbers of other taxa	7	9	1276.88 $\pm$ 604.34
	mostly open substrate with patches of low density sea pens, hydroids and ascidians with moderate numbers of individual echinoids	8	2	321.53 $\pm$ 76.96
Low/Medium density benthic community	mostly open substrate with patches of medium density bivalves & low numbers of individual polychaetes	10	1	5.09 $\pm$ 0.48
	mostly open substrate with patches of low/medium density hard and soft coral with sea pens, and low numbers of individual echinoids and bivalves	12	1	90.33 $\pm$ 50
Medium density benthic community	patches of medium density bivalves & low numbers of individual polychaetes, with areas of open substrate	9 & 11	2	11.83 $\pm$ 1.41
	patches of medium density sea pens, hydroids & ascidians with areas of open substrate and low numbers of individual bryozoans, soft coral, echinoids, crustaceans & bivalves	13	10	1454.56 $\pm$ 837.26
<b>Total</b>		<b>13</b>	<b>572</b>	<b>6898.29 <math>\pm</math> 3863.60</b>

Figure B7.4.6.2f Comparison of Algae Community Regions at Cairns Harbour over time, 2001 and 2012/13 (McKenna et al. et al. 2013)





### B7.4.6.3 Infauna Communities

For macroinvertebrate infauna during the dry season, indices of taxonomic richness, abundance, and Shannon's diversity were highest at the offshore sites 5C and 5D, found within sediment class 10 (refer **Appendix D13, Figure D13.4d**). Some sites within sediment classes five and seven tended to have lower abundance, richness and diversity than surrounding sites, but these low values were not consistent within either sediment classes. Diversity measures at the existing spoil ground, represented by sites 7A, 7B, and 7C were within the range of variation observed at surrounding sites. During the wet season, diversity indices were lower at all sites, however, variation in these indices was similar across sites and sediment classes with greatest richness, abundance and Shannon's diversity seen in sediment classes one, five and 10. Once again, diversity measures at the existing spoil ground were within the range of variation observed at surrounding sites.

Higher abundance and richness within sediment class 10 reflected the large numbers of polychaetes and crustaceans (**Figure B7.4.6.3a**) from numerous taxonomic groups. Although richness and abundance was lower at these same sites during the wet season, the taxonomic distribution at these sites remained similar, with crustaceans and polychaetes dominating. Higher richness and abundance at site 4A during the dry season, where large numbers of crustacean species were recorded, may be associated with local hydrodynamics, as this site was located at the edge of the outer shipping channel. It should be noted that site 1A from sediment class 11 was taken from Trinity Inlet and is unlikely to be representative of this sediment class in the offshore area, where it was predominantly found.

Sea-pen 1 (**Figure B7.4.6.2a-A**) was large enough to be observed in video transects as a part of the epibenthos, and small enough to be captured within van Veen grabs with other infauna. It was particularly abundant around the existing DMPA (at sites 7A, 7B, 7C, 8A and 8B) and was seen in both surveys, although it was most abundant during the dry season. These results were consistent with the results of McKenna *et al.* (2013), where these small sea pens were major community components surrounding the existing DMPA.

Infaunal communities from the proposed DMPA, taken in July 2014, had similar species distributions to other sites in the area with crustaceans and polychaetes dominant (**Figure B7.4.6.3b**).

Figure B7.4.6.3a Proportion of each Major Taxon Contributing to Total Abundance at each Site during Wet Season and Dry Season Surveys





Figure B7.4.6.3b Proportion of each Major Taxon Contributing to Total Abundance at DMPA Sites



Multivariate analysis of dry season data showed that infaunal communities varied among acoustically derived substrate classes (ANOSIM global  $R=0.28$ ,  $p=0.002$ ). Communities from sediment class five occupied a central position within the plot and were very similar to a range of communities (not significantly different) including those from sediment classes one, 9 and 11. The most disparate communities came from sediment class eight, which were very different from communities in all other sediment classes except from class six. Typically, similar substrate types tended to have similar communities; sediment classes (e.g. one and five were similar, five and six were similar, and six and seven were similar, etc.). The only adjacent sediment classes with different communities were classes seven and eight, and classes nine and 10. Because sediment classes seven and eight surrounded the existing DMPA, the potential effect of the existing DMPA was also explored in more detail.

While sediment class was a reasonable predictor of infaunal macroinvertebrate communities, location (with respect to coastline and depth) had better explanatory power (**Appendix D13**) (ANOSIM global  $R=0.664$ ,  $p=0.0001$ ). The inner channel consisted of channel sites inside Trinity Inlet, the outer channel was channel sites in the Cairns harbour, coastal sites were close to the shore and within the one metre LAT contour, inshore sites were between the five-10 m LAT contours, mid-shore sites were within 10-20 m LAT contours, and offshore sites were deeper than 20 m LAT. Communities from all locations were significantly different to one another ( $p<0.002$ ). The inner channel community was dominated by polychaetes (glycerids, maldanids, and capitellids) and tellnid bivalves, the outer channel was characterised by tellnids and amphipod crustaceans. The coastal sites were characterised by large numbers of marine worms (spionids, orbininids, nemerteans) and amphipods, inshore sites were dominated by amphipods, mid-shore sites by amphipods and sea pens, while the offshore sites had large numbers of crustaceans (tanaids, amphipods, callianassids ghost shrimp) and maldanid polychaetes.

Communities at the existing and proposed DMPAs clustered centrally with respect to surrounding background sites (**Appendix D13**). This ordination shows the spoil options have communities that are more similar to the existing DMPA than at surrounding background and references sites.

An n-MDS ordination of the spoil options and surrounding references sites (5A,5B, 5C) shows that the sites closest to the existing DMPA in geographical space, are also the most similar in community space (**Appendix D13**). That is, communities at Option 1 are the most similar to the existing DMPA communities, and these are also the two closest locations. Reference sites 5c and 5d are located in substantially deeper water and these clustered together to the left of the plot. While the different spoil option communities were different to one another (ANOSIM global  $R=0.619$ ,  $p=0.0001$ ), the magnitude of these differences were minor in comparison with gross community differences across the entire data set.

In summary, infaunal communities differed among broad habitat types, but these differences could not consistently be determined among all of the acoustically-derived substrate classes. Physical location was a better community predictor, particularly when sites were grouped according to distance offshore and water depth. The existing DMPA communities were most similar to the northern adjacent sites. This may be related to the northerly movement of placed dredged material, driven predominantly by south-east winds. Alternatively, it may simply be related to proximity as relative differences in community space and geographic space were similar. All of these trends were reasonably consistent between wet and dry seasons, albeit with greater variation between sites in the wet season than during the dry season.

### **B7.4.7 Marine Species of Conservation Significance**

This description of marine species of conservation significance is based on a desktop review of existing information, together with advice acquired through liaison with local experts. In agreement with the relevant regulators, a targeted marine megafauna survey was not deemed necessary for this EIS on the basis that the proposal does not involve seabed reclamation or direct loss of seagrass, applicable mitigations measures are well understood, and that surveys will only provide a limited 'snapshot' for any given time. Further, any present surveys would likely grossly under-estimate typical population estimates and habitat utilisation for seagrass-dependent species (e.g. dugongs, turtles), given the limited availability and poor condition of local seagrass communities at this time.

While there is very limited data available describing the occurrence, habitat utilisation and populations of marine megafauna in the Cairns region, a conservative approach has been adopted, whereby it has been assumed that a species may occur i) if suitable habitat is available, and ii) the area of concern is within its broader geographical range.

The EPBC Act Protected Matters Search Tool was used to identify threatened marine species and ecological communities that occur or could occur within the study area. In summary, the following were identified:

- Threatened sharks: three species
- Threatened marine mammals: two species
- Threatened marine reptiles: six species
- No threatened ecological communities.

An additional seven species were listed as protected migratory species, including five mammals, one reptile and one shark.

Further to the threatened and migratory species, numerous species were listed only as 'listed marine species' (i.e. non-threatened, non-migratory). These additional EPBC Act listed marine species are not addressed in detail in this section but, for future reference, included 50 syngnathids (i.e. seahorses, pipehorses, pipefish), 15 sea snakes, and six mammals (minke whale and five dolphins).

Detail is provided for key fauna groups in the following sections.

#### **B7.4.7.1 Marine Turtles**

Six species of marine turtle may occur in the harbour, Trinity Inlet and/or the surrounding area, although some species are more common than others (**Table B7.4.7.1a**). All are considered threatened under the EPBC Act and NC Act. Marine turtles have been recorded in offshore, near-shore and intertidal habitats within the study area. These species primarily use the area as a feeding ground, as well as a transit area between other parts of the Queensland coast and the Great Barrier Reef. Low density nesting also occurs across the broader region, particularly low density nesting of Green and Flatback turtles (Limpus 2007, 2008).

Table B7.4.7.1a EPBC Protected Matters Database Search Results – Threatened and Migratory Marine Reptiles

Species	Status (EPBC/ NCA)*	Distribution / Habitat	Study Area (Cairns Harbour)	Offshore DMPA(s)
<i>Caretta caretta</i> Loggerhead Turtle	EPBC: E, M, LM NCA: E	Pelagic and benthic species. Cairns region is not known to represent a nesting site. Forages on marine invertebrates (Wilson and Swan 2004).	Likely – foraging habitat available at seagrass areas and coastal rocky shores.	Likely - suitable habitat located at surrounding reef areas.
<i>Chelonia mydas</i> Green Turtle	EPBC: V, M, LM NCA: V	Marine waters and near the seabed. Foraging habitat available at seagrass meadows, where it feeds mainly on seagrass and benthic invertebrates (Wilson and Swan 2004). Low density nesting occurs in broader region, with known hotspots at the sandy beaches between Cairns and Hinchinbrook (e.g. Cowley and Kurrimine Beaches).	Likely – strong association with seagrass (noting low seagrass occurrence at present), reefs (algae) and mangroves.	Likely - suitable habitat located at nearby reef and seagrass areas.
<i>Dermochelys coriacea</i> Leathery Turtle, Leatherback Turtle	EPBC: V, M, LM NCA: E	Oceanic species, feeds on jellyfish and other soft bodied invertebrates (DEWHA 2007, Wilson 2005). Rarely sighted in Cairns region and then only in deep waters. Rarely nests on northern Queensland coastline.	Unlikely – oceanic species.	Possible transient visitor.
<i>Eretmochelys imbricata</i> Hawksbill Turtle	EPBC: V, M, LM NCA: V	Forages on coral and rocky reefs, sponges major dietary component. Cairns region not a known nesting area (normally occurs north of Princess Charlotte Bay).	Possible transient visitor, some foraging.	Possible transient visitor.
<i>Lepidochelys olivacea</i> Olive Ridley Turtle, Pacific Ridley Turtle	EPBC: E, M, LM NCA: E	Marine waters over soft sediments, but not common. No known nesting in Cairns region.	Unlikely – oceanic species	Possible - uncommon transient visitor
<i>Natator depressus</i> Flatback Turtle	EPBC: V, M, LM NCA: V	Marine species found around reef areas and soft sediment habitats; forages on invertebrates. Primarily nests south of Mackay, though low density nesting (<10 females per year) in Cairns region.	Likely – suitable habitat and common within GBR lagoon.	Likely – suitable habitat and common within GBR lagoon.
<i>Crocodylus porosus</i> Saltwater Crocodile	EPBC: M, LM NCA: V	Coastal waters, estuaries and reefs throughout geographic range, including Cairns region.	Likely – throughout study area	Possible – more common inshore

(\* ) E = endangered; V = vulnerable; M = migratory; NT = near threatened; LM = listed marine

Overall, the distribution and abundance patterns of turtles within the study and project areas are thought to be mainly a function of the availability of suitable food resources, as summarised below:


- Green turtle (*Chelonia mydas*) is listed as vulnerable under both the EPBC Act and the NC Act. It feeds directly on seagrasses and algae (Brand-Gardner et al. 1999) with most foraging activity concentrated around the seagrass beds of Cairns harbour and Trinity Inlet. Within the study area, a low incidence of nesting occurs along the northern beaches, tending to concentrate at Clifton Beach (pers. comm. J. Gilbert). There are also anecdotal reports of nesting at the northern end of the Cairns Esplanade. Beyond this, green turtle nesting has occasionally been observed at Green Island, Fitzroy Island and Michaelmas Cay. Green turtles are historically the most common marine turtle species in the study area but in recent years, since Tropical Cyclone Yasi in 2007, there has been increased strandings and mortality for this species (refer **Figure B7.4.7.1a** for recent marine turtle stranding summary). The cyclone resulted in a drastic reduction in seagrass extent and condition, causing reduced food availability, which led to turtle starvation, mortality and reduced fecundity. The local population is likely to be subject to ongoing stress until significant seagrass recovery occurs
- Loggerhead turtle (*Caretta caretta*) is listed as endangered under both the EPBC Act and the NC Act. It is a carnivorous species which feeds on jellyfish, crustaceans, echinoderms, and bivalve molluscs from seagrass meadows and reef areas (Limpus et al. 1994). Given the current reduced seagrass availability at present, this species is likely to be more abundant around the offshore reefs of the surrounding area, but may also potentially forage around the coastal rocky shores east of the Cairns harbour
- Hawksbill turtles (*Eretmochelys imbricata*) are listed as vulnerable under both the EPBC Act and the NC Act. They feed primarily on sponges, but also feed on seagrasses, algae, soft corals and shellfish. Hawksbill turtles would most commonly occur around the offshore reefs of the surrounding area where sponges are likely to be abundant, but they may also potentially forage around the coastal rocky shores east of the Cairns harbour. Similar to green turtles, there have also been reports of hawksbill turtles suffering starvation in the Cairns area in recent times (pers. comm. J. Gilbert)
- Flatback turtles (*Natator depressus*) are listed as vulnerable under both the EPBC Act and the NC Act. They are carnivorous, feeding mainly on soft-bodied invertebrates (sea cucumbers, sponges, soft corals, jellyfish, etc.) from the sea floor (Wilson 2005; Wilson and Swan 2003)
- Olive Ridley turtles (*Lepidochelys olivacea*) are listed as endangered under both the EPBC Act and the NC Act. They are thought to be uncommon, occurring in very low numbers throughout the region and preferring waters over soft sediments. This species is thought to be mostly carnivorous, with a diet mostly comprised of urchins, small crabs, and molluscs (Wilson 2005)
- Leatherback turtles (*Dermochelys coriacea*) are listed as vulnerable under the EPBC Act and endangered under the NC Act. They are uncommon in the local area, typically occurring in deeper oceanic waters. Leatherbacks feed on jellyfish and other soft bodied invertebrates (DEWHA 2007; Wilson 2005)

With the exception of green turtles, there is a lack of data to describe key or preferred foraging habitats for the remaining marine turtles in the Cairns region, with the less common species (i.e. small resident populations) scarcely being studied nation-wide. It is likely, however, that marine turtles that exist within the study area and DMPA project areas are transients, rather than residents. This would primarily be due to the lack of i) optimal or perennial feeding resources in this area, and ii) suitable nesting habitat (sandy beaches). However, it is likely that the sparse seagrass and epibenthic fauna assemblages in the study area are used sporadically or occasionally by some marine turtles. Loggerhead turtles may also feed on jellyfish that occur in the pelagic waters of the study area.

While abovementioned stranding data refers to total marine turtles, the most recent available StrandNet database annual report (Meager and Limpus 2012) provides species-specific records of three turtle species confirmed stranded in the Cairns region (latitudinal grids 16 and 17) during 2011, and hence using the study area and surrounding waters relatively recently. These included green turtle (66), hawksbill (10) and flatback (1), with green turtle being by far the most commonly stranded at this time. Note that Hazel and Gyuris (2006) identified Cairns as one of five Queensland hot-spot areas for vessel-related sea turtle strandings based on analyses of StrandNet data (i.e. concentrated strandings).

The coastline of the study area is not a critical turtle breeding area, with most turtles residing in the study believed to have originated from rookeries elsewhere on the central and Far North Queensland coast and islands. The exceptions to this are green and flatback turtles. While low density nesting for these species may occur on sandy beaches within the study area (e.g. Yorkeys Knob), the majority of marine turtle nesting within the region occurs south of Cairns on the sandy beaches towards Hinchinbrook, and further north at Princess Charlotte Bay (Limpus 2007, 2008).

**Figure B7.4.7.1a Number Marine Turtle Strandings Recorded in the Queensland Strandings Database for 1 January – 30 June 2013, with Comparison to the same Period for Previous Years (Department of Environment and Heritage Protection 2013)**



### B7.4.7.2 Other Marine Reptiles

Saltwater crocodile *Crocodylus porosus* is a listed migratory and marine species under the EPBC Act, and is considered a vulnerable threatened species under the NC Act. Saltwater crocodiles occur in the study area and surrounds, with multiple records within the study area registered on the Wildlife Online database (DSITIA 2013). Nesting sites and preferred feeding habitats in the study would likely be restricted to Trinity Inlet (i.e. typically prefers mangrove-lined estuaries).

The EPBC Protected Matters database search results indicate that 15 species of sea snake occur or could occur in the study area. These species are listed marine species and are protected under the EPBC Act, but are not considered to be threatened under Commonwealth or Queensland legislation.

### B7.4.7.3 Threatened Sharks

Three threatened shark species were identified as potentially occurring in the study area using EPBC Protected Matters search tool: whale shark *Rhincodon typus*, dwarf sawfish *Pristis clavata* and green sawfish *Pristis zijsron* (Table B7.4.7.3a). The whale shark is a pelagic species that tends to prefer offshore tropical waters. There are occasional records of this species in the surrounding area, although it is considered to represent a transient visitor to the region. Green sawfish and dwarf sawfish may occur in the study area from time to time, with Cairns representing the present day southern extent of their geographic distribution along the Queensland coast. Based on data from museums, research surveys and fisheries observations, Stevens et al. (2005) report that green sawfish have been recorded in the Cairns area in recent decades (i.e. since 1990). However, long-term data from the Queensland Shark Control Program indicate a major population decline has occurred since the 1970s, largely due to fishing pressure (targeted and bycatch) and net entanglement. While both green and dwarf sawfish may occur within the study area, it is likely that the local population is very small and/or transient. Dwarf sawfish are typically considered to be the more common *Pristis* species throughout northern Australia (Stevens et al. 2005), although there are no known recent records from east of Cape York (GBRMPA 2012).

While not a threatened species, the Porbeagle Shark *Lamna nasus* is listed under the EPBC Act as a migratory marine species. This shark is unlikely to occur within the study area or surrounding waters as it is typically associated with oceanic waters offshore and along the east coast it is thought to be limited to southern Queensland and south-eastern Australia.

The EPBC Protected Matters database search results indicate that 50 species of sygnathids (seahorse, pipe horse and pipefish) occur or could occur in the study area. These species are listed marine species and are protected under the EPBC Act, but are not considered to be threatened under EPBC or state legislation.

Table B7.4.7.3a EPBC Protected Matters Database Search Results – Threatened and Migratory Shark Species

Species	Status (EPBC/ NCA)*	Distribution / Habitat	Study Area (Cairns Harbour)	Offshore DMPA(s)
<i>Pristis zijsron</i> Green Sawfish, Dindagubba, Narrowsnout Sawfish	EPBC: V	Cairns thought to represent extent of southern distribution. Occurs in estuaries and river mouths, embankments and beaches. While more common nearshore, has been recorded offshore. Benthic feeder, found in to 70 m (TSSC 2008).	Likely, though may be small or transient population.	Possible - uncommon transient visitor.
<i>Pristis clavata</i> Dwarf Sawfish, Queensland Sawfish	EPBC: V	Cairns thought to represent extent of southern distribution. Occurs nearshore in estuaries and river mouths, embankments and beaches. Benthic feeder, found in depths from 1 m to 10 m (TSSC 2008).	Likely, though may be small or transient population.	Unlikely, near shore species.
<i>Rhincodon typus</i> Whale Shark	EPBC: V, M	Wide ranging tropical species. Critical habitat in Australia includes Ningaloo Reef in WA, the Coral Sea and Christmas Island. Cairns region not known to represent an important habitat for this species.	Unlikely – Low abundance regionally and lack of deep waters limit habitat value of the study area.	Possible – uncommon transient.
<i>Lamna nasus</i> Porbeagle, Mackerel Shark	EPBC: M	Pelagic and oceanic waters offshore, though likely limited to the cooler waters of southern Queensland`.	Unlikely – oceanic species and study area beyond known geographic distribution.	Unlikely – oceanic species and DMPAs beyond known geographic distribution.

(\*) E = endangered; V = vulnerable; M = migratory; NT = near threatened; LM = listed marine

#### B7.4.7.4 Marine Mammals

There are seven listed migratory marine mammals that could occur within or adjacent to the study area (**Table B7.4.7.4a**). Five are considered to be threatened, or near threatened, under the EPBC Act and/or NC Act. Of the latter, the four marine mammal species with a likely occurrence in the study area and the project areas are the Australian snubfin dolphin (*Orcaella heinsohni*), Indo-Pacific humpback dolphin (*Sousa chinensis*), dugong (*Dugong dugon*) and humpback whale (*Megaptera novaeangliae*). These four species are discussed in further detail below.



Table B7.4.7.4a EPBC Protected Matters Database Search Results – Threatened and Migratory Marine Mammals

Species	Status (EPBC/ NCA)*	Distribution / Habitat	Study Area (Cairns Harbour)	Offshore DMPA(s)
<i>Balaenoptera musculus</i> Blue Whale	EPBC: E, M, LM	Oceanic waters	Unlikely – shallow water depths and port infrastructure limits values.	Unlikely – transient offshore.
<i>Megaptera novaeangliae</i> Humpback Whale	EPBC: V, M, LM NCA: V	Oceanic waters	Unlikely – shallow water depths and port infrastructure limits values.	Possible - Breeding habitat.
<i>Balaenoptera edeni</i> Bryde's Whale	EPBC: M LM	Oceanic waters off Australian and southern Africa where it searches for baitfish (Van Dyck and Strahan 2008).	Unlikely – shallow water depths and port infrastructure limits values.	Unlikely – transient offshore.
<i>Orcaella heinsohni</i> (previously <i>O. brevirostris</i> ) Australian Snubfin, Irrawaddy Dolphin	EPBC: M, LM NCA: NT	Shallow coastal and estuarine waters.	Possible – Past records nearby around Ellis Beach area.	Possible – primarily an inshore species.
<i>Orcinus orca</i> Killer Whale, Orca	EPBC: M, LM	Oceanic, pelagic and relatively shallow waters over continental shelf. Marine mammals are key dietary component (Van Dyck and Strahan 2008).	Unlikely – shallow water depths and port infrastructure limits values.	Possible – may occur offshore waters (in part associated with Humpback Whale breeding)
<i>Sousa chinensis</i> Indo-Pacific Humpback Dolphin	EPBC: M, LM NCA: NT  EPBC: M, LM NCA: V	Shallow coastal and estuarine waters preferred; recorded in offshore waters within GBR lagoon (i.e. sheltered by reef).	Possible – preferred habitat available, past records nearby around Ella Bay.	Possible – though primarily an inshore species.
<i>Dugong dugon</i> Dugong		Inshore coastal and estuarine waters where notable seagrass meadows occur.	Likely – strong association with seagrass (noting low seagrass occurrence at present).	Possible – transient.

(\* ) E = endangered; V = vulnerable; M = migratory; NT = near threatened; LM = listed marine

## Dolphins

The two threatened dolphin species likely to occur, Snub fin and Indo-Pacific humpback dolphins, are considered a high priority by GBRMPA as numbers of these inshore species are thought to be in decline. Key threats driving this population decline are boating interactions, net entanglement and poor water quality, which can affect prey resources.

The Australian snubfin dolphin has a global IUCN listing of “near threatened” (IUCN 2010). It is listed as a migratory marine mammal under the EPBC Act and near threatened under the NC Act. Studies to date indicate that this species generally occurs in waters less than 15 m deep, within 10 km of the coast and within 20 km of a river mouth (Parra et al. 2004). The species is an opportunistic generalist, feeding on fish and cephalopods (octopus, squid etc.) from coastal, estuarine and near-shore reef habitats (Parra 2006, Parra et al. 2006b). Meager et al. (2012) state that the only reliable population estimate for this species is from Cleveland Bay (Townsville), where the sub-population had been estimated between 64 and 76 individuals in 2000-2001.

Indo-Pacific humpback dolphin has a global IUCN listing of “near threatened” (IUCN 2010). It is listed as a migratory marine mammal under the EPBC Act and near threatened under the NC Act. In Australia its distribution stretches from northern New South Wales along the coast of Queensland to Shark Bay in Western Australia. Studies to date indicate that this species like the Australian snub-nosed dolphin generally occurs in waters less than 15 m deep, within 10 km of the coast and within 20 km of a river mouth (Parra et al. 2004). The species is also an opportunistic generalist, feeding on fish and crustaceans from coastal, estuarine and near-shore reef habitats (Parra 2006, Parra et al. 2006b). No reliable population estimates are available for northern Queensland regions (Meager et al. 2012).

Detailed marine mammal surveys undertaken for a port elsewhere in Queensland where these species commonly occur (Port of Gladstone: Parra 2006, GHD 2009) found that the Indo-Pacific humpback dolphin had important feeding and nursing areas in the vicinity of estuary/river mouths, and favoured shallow waters (one-two metres deep) where seagrass is present. In contrast, the snubfin dolphin favoured water two to five metres deep in dredged channels. If similar preferences are transferable to the present study area, it is likely that these species would prefer the shallow waters of Cairns harbour and the mouth of Trinity Inlet.

*Sousa chinensis* is occasionally recorded from the Cairns area in low numbers on the StrandNet database, with a total of four strandings from 2008 to 2010. Indo-Pacific Humpback dolphins have recently been sighted in the project area, including two individuals at the marina in Trinity Inlet in November 2013 and two individuals swimming in shallow waters off Ellis Beach in September 2013 (personal observations: Ms Emma Scott). There are no known recent sightings of snubfin dolphins in the project area. However, this species is known to regularly frequent the Port Douglas and Low Isles areas, approximately 35 km north of the project area, typically in groups of up to six individuals (personal observations: Ms Emma Scott). Given declining numbers and small population sizes, it is thought that these two threatened dolphin species can only sustain very small levels of anthropogenic mortality (Meager et al. 2012).


## Dugong

The dugong has a global IUCN listing of “vulnerable to extinction” (IUCN 2010). It is ‘listed threatened’, ‘listed migratory’ and ‘listed marine’ species under the EPBC Act and the Queensland dugong population is considered “vulnerable” under the NC Act. Dugongs are principally herbivores and have been shown to be highly selective feeders, preferring certain species of seagrass to others. Dugongs have a preference for *Halophila* spp., which can be prevalent in seagrass meadows within the study area and surrounding waters, particularly the Cairns harbour and Trinity Inlet, and those areas along the eastern shoreline toward Bessie Point and Cape Grafton.


Most broad scale dugong surveys and population estimates have focused on the southern Great Barrier Reef, with limited coverage of the Cairns area (e.g. Marsh and Saalfeld 1990, Marsh et al. 1999). Marsh et al. (1996) also incorporated analysis of dugong records from historical shark net captures (i.e. Queensland Shark Control Program nets), whereby 101 dugongs were killed in shark nets off Cairns between 1962 and 1978. Overall, the general consensus of these studies is that there has been a strong decline in dugong catches since the 1960s, to the point that the sightings are now too few to enable a population estimate. As far as local distribution, targeted surveys by Marsh and Saalfeld (1990) confirmed that the dugongs recorded in the Cairns area were sighted close to inshore seagrass beds.

While dugongs can be common in Trinity Bay, it is thought that current population numbers are low due to the reduction in the extent and condition of local seagrass meadows. The year 2011 saw record dugong strandings throughout northern Queensland (Meager and Limpus 2012b; **Figure B7.4.7.4a**), following the depletion of cyclone-affected seagrass food resources. Since then, it is likely that the majority of residents would have had to forage elsewhere in Queensland or Torres Strait to avoid malnutrition and starvation in the local area. A combination of emigration and past mortality is likely responsible for the reduced stranding records over the last two years (**Figure B7.4.7.4b**), and may be considered a proxy for a reduction in the local population size as compared with previous years.

**Figure B7.4.7.4a Monthly Dugong Strandings in Northern Queensland during 2011, Compared with Long-Term Average (1996-2012) (from Meager and Limpus 2012b)**



**Figure B7.4.7.4b Number of Dugong Strandings Recorded in the Queensland Strandings Database for 1 January – 30 June 2013, with Comparison to the same Period for Previous Years (DEHP 2013)**




## Humpback Whale

The humpback whale has a global IUCN listing of least concern (IUCN 2010). It is listed as vulnerable under both the EPBC Act and the NC Act. Humpback whales calve in the protected waters of the GBR between July and August then travel along the Australian coast to Antarctic waters where they spend spring and summer before returning (DERM 2010). They feed on krill (*Euphausia superba*) and small fish.

The waters surrounding Cairns have been modelled to be of moderate environmental suitability for humpback whales compared to the highly suitable waters offshore from Mackay (Figure B7.4.7.4c). Although the area off Cairns is comparably less suitable to other areas of the GBR lagoon, large numbers of whales are sighted off Cairns each winter. The regularity and increasing trend of sightings has led to the commencement of whale watching operations out of Cairns.

**Figure B7.4.7.4c Model Prediction of Environmental Suitability for Humpback Whales in the GBRWHA**  
(Source: Smith et al. 2012). Red = Most Suitable; Green = Moderate Suitability; Blue = Least Suitable)



## B7.4.8 Exotic Marine Species and Pests

More than 250 non-indigenous marine species have been recorded in Australian waters (NIMPCG 2011). There are several potential vectors by which non-indigenous species may enter domestic waters; however it is thought that most species are unintentionally introduced through shipping and vessel movements, either in ballast waters or from biofouling on the hull of vessels (Hewitt and Campbell 2010). Other vectors include intentional transfer of aquaculture and mariculture organisms, transfer of food products for the aquarium trade and use of biological material for packing (Hewitt and Campbell 2010).

There are currently no known marine pests established in the Cairns area (Australian Government 2013). Risk tables, developed as part of the Australian Ballast Water Management Information System (ABWMIS), identify species of known or potential concern for Australian ports. The 2013 risk tables identify Cairns as a potential recipient port for the Asian date mussel *Musculista senhousia* and European fan worm *Sabella spallanzani*. At present, *S. spallanzanii* has not previously been recorded in Cairns but is known to occur as a pest at southern Australian ports in temperate waters. *M. senhousia* was recorded fouling a Navy vessel at Cairns in 2007 (refer below) but has not been detected here since by either the Queensland Government, or though Ports North's routine marine pest monitoring activities.

Other introduced species have previously been identified from the Port of Cairns, including Asian green mussel *Perna viridis* and Caribbean tubeworm *Hydroides sanctaecrucis* (Environment North 2005, Souter 2009). While these species pose a potential threat, neither are considered to be pest species currently established in Australian waters. Both species are prolific fouling organisms with the potential for causing extensive nuisance growths. In 2001, government surveys recorded *Perna viridis* at Cairns in low numbers on both vessel hull(s) and in Trinity Inlet, with the variety of age classes present suggesting that successful local spawning had occurred. During the same year, *Hydroides sanctaecrucis* was recorded on hard substrata at the Port of Cairns, particularly between the Marlin Marina and Smith's Creek (Environment North 2005). Following the discovery of these species at Cairns in 2001, various eradication measures were implemented with neither species being detected during subsequent 2002 surveys (Environment North 2005).

In 2007 an outbreak of both *Perna viridis* and *Musculista senhousia* (Asian green mussel and Asian date mussel) were recorded fouling a vessel at Cairns, with *Perna viridis* again being recorded on a vessel in 2008 (Australian Government 2013). Appropriate control measures were implemented with subsequent monitoring undertaken following an initial quarantine period. The Australian Marine Pests website has not reported any further marine pest outbreaks for the Cairns region.

## B7.4.9 Fish and Shellfish Resources

### B7.4.9.1 Fish Communities

Coles et al. (1993) undertook intensive sampling of fish communities within Cairns Harbour quarterly in 1988. Sites were positioned in seagrass meadows, and sampled using beam trawls, gill nets and seine nets. In total, 5614 fish were collected, comprised of 134 fish taxa. Of the 134 species recorded, only five were considered to be of high commercial or recreational fisheries value. The most numerous fish species were two small-bodied demersal species: a goby (*Yongeichthys criniger*) and a ponyfish (*Leiognathus splendens*). The highest biomass was contributed by larger-bodied species, namely *Carcharhinus* sharks, salmon *Polydactylus sheridani* and *Eleutheronema tetradactylum*, the queenfish *Scomberoides commersonianus* and the catfish *Arius proximus*. Very few reef fish species were recorded; most species are pelagic or soft sediment-associated species.

Coles et al. (1993) found that small-bodied fish and juveniles dominated on seagrass meadows, and were found in similar abundance (and size range) reported between Cairns and Bowen, and at Mornington Island. This is consistent with case studies elsewhere (e.g. Carruthers et al. 2002; Bell and Pollard 1989) which demonstrate the importance of seagrass meadows as a fish nursery habitat. The presence of larger predators also highlights the importance of seagrass meadows as feeding habitats for fish.


Blaber (1980) surveyed the fish fauna of Trinity Inlet and unvegetated tidal sand banks within Cairns Harbour using seine nets, gill nets and stake nets. He found that unvegetated flats in Cairns Harbour had (i) fewer juveniles; (ii) higher numbers of piscivores (mostly adults); and (iii) fewer planktivores than sites in Trinity Inlet. On this basis, Blaber (1980) argued that mangrove-lined creeks within Trinity Inlet represented higher value nursery habitats for fish than those found in Cairns Harbour.

These two studies both demonstrate the importance of both mangrove-lined estuaries and seagrass meadows within the study area as nursery and adult feeding habitats. Unvegetated soft sediment habitats also represent important habitats for several species of commercial and recreational significance. For example, species of fisheries importance such as grunter/javelin fish (*Pomadasys kaakan*), nannygai (*Lutjanus malabaricus*, *L. erythropterus*), flathead all use unvegetated soft sediment habitat during one or more parts of their life-cycle (Table B7.4.9.1b). Note that the fish species thought to dominate local recreational catches, coral trout and snapper (Figure B7.4.9.1a), are typically reef-associated species.

Building on the above baseline surveys, the Queensland Department of Primary Industries annual assessments of stocks of key fisheries species at Trinity Inlet and the Cairns harbour from 1997 to 2000, which targeted barramundi, king threadfin, blue threadfin, mangrove jack, grunter, bream and mud crab (Helmke et al. 2003; Figure B7.4.9.1a). While not all of these target species were captured, the surveys did record a total of 462 fish representing 65 species from 29 families. The most frequently caught species varied between both survey years and sites. Overall, barramundi and saltwater bony bream were the most commonly caught in 1997, compared with anchovy in 1998. The year 2000 recorded relatively low catches across all common species. In terms of differences in the distribution catches across the sites surveyed by Helmke et al. (2003), notably high catch rates included those for anchovy, saltwater bony bream and Burnett salmon in the main channel of Trinity Inlet, and barramundi at Saltwater/Little Barron River (Table B7.4.9.1a)

The fish communities of the study area show great variability over a range of temporal scales. At seasonal time scales, many fish species found in the study area have a peak spawning and/or recruitment period in late spring-summer (Kailola et al. 1993; Table B7.4.9.1b). Many estuarine fish also show great variability in abundance in response to cyclic feeding patterns (diel, tidal and lunar patterns), breeding and spawning activities, and in response to rainfall and other environmental processes.

**Figure B7.4.9.1a Catch Rates of the Most Frequently Caught Fish from Surveys at Trinity Inlet and Cairns Harbour, 1997, 1998, 2000 (from Helmke et al. 2003)**



**Table B7.4.9.1a Total Catches of the Most Frequently Caught Fish from Surveys at Individual Sites Throughout Trinity Inlet and Cairns Harbour, 1997, 1998, 2000 (from Helmke et al. 2003)**

Family	Common Name	Year	Total Number	False Cape	Catch Rate (number per hour)						
					Bessie Point	Saltwater - Little Barron	Main Channel	Smiths Creek	Redbank Creek	Skeleton Creek	
Engraulidae	Hamilton's anchovy	1997	5	0.00	*	0.50	0.28	0.00	0.00	0.00	0.08
		1998	62	0.00	0.27	0.41	3.16	0.81	0.00	0.00	0.07
		2000	11	0.00	0.00	0.00	*	0.30	0.00	0.00	0.76
Centropomidae	barramundi	1997	29	0.00	*	6.25	0.00	0.08	0.20	0.00	0.00
		1998	27	0.12	0.55	0.90	0.00	0.16	0.18	0.00	0.15
		2000	5	0.00	0.05	0.09	*	0.07	*	*	0.00
Clupeidae	salt water bony bream	1997	26	0.00	*	0.50	0.00	0.50	0.00	0.00	0.08
		1998	8	0.00	0.00	0.00	2.42	0.24	0.00	0.00	0.39
		2000	5	0.00	0.00	0.00	*	0.23	*	*	0.10
Mugilidae	mullet	1997	13	0.25	*	0.25	0.00	0.66	0.00	0.00	0.16
		1998	5	0.27	0.09	0.08	0.00	0.00	0.00	0.00	0.00
		2000	9	0.00	0.00	0.00	*	0.69	*	*	0.00
Polynemidae	Burnett salmon	1997	0	0.00	*	0.00	0.00	0.00	0.00	0.00	0.00
		1998	18	0.18	0.09	3.39	0.00	0.00	0.00	0.00	0.00
		2000	4	0.00	0.05	0.47	*	0.00	*	*	0.00
Clupeidae	deep herring	1997	15	0.00	*	0.00	0.00	0.00	0.00	0.00	1.25
		1998	0	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		2000	0	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Scaenidae	silver jewfish	1997	3	0.00	*	0.00	0.28	0.00	0.00	0.10	0.00
		1998	3	0.00	0.09	0.22	0.07	0.00	0.00	0.00	0.00
		2000	9	0.00	0.05	0.36	*	0.00	*	*	0.00
Clupeidae	herring	1997	13	0.00	*	0.00	0.00	0.40	0.00	0.00	0.75
		1998	0	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		2000	0	0.00	0.00	0.00	*	0.00	*	*	0.00
Mugilidae	diamond scaled mullet	1997	9	0.87	*	0.00	0.14	0.00	0.00	0.00	0.08
		1998	1	0.00	0.00	0.08	0.00	0.00	0.00	0.00	0.00
		2000	3	0.00	0.59	0.00	*	0.00	*	*	0.00
Ariidae	catfish	1997	3	0.35	*	0.00	0.00	0.00	0.00	0.00	0.00
		1998	0	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		2000	10	0.00	0.59	0.00	*	0.00	*	*	0.00

Table B7.4.9.1b Key Fisheries Species Present in the Study Area and their Primary Habitats at Different Stages of their Life-Cycle (Data: mostly Kailola et al. 1993)

Common Name	Species name	Spawning period	Estuary						Estuary									
			Mangroves	Seagrass	Shoals	Channels/ deeper soft sediment	Freshwater	Inshore sediment*	Offshore sediment	Open water	Reefs*							
<b>Fish</b>	<b>Teleosti</b>																	
Threadfin salmon	<i>Polydactylus macrochir</i> , <i>Eleutheronema tetradactylum</i>	Oct-Apr (peak Dec)	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.											
Barramundi	<i>Lates calcarifer</i>	Oct-Mar (peak Dec)	Spw., Juv, Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.									
Dusky flathead	<i>Platycephalus fuscus</i>	Sep-Mar	Juv., Ad.	Spw., Juv., Ad.	Spw., Juv., Ad.	Ad., Juv.			Spw.									
Sand whiting	<i>Sillago ciliata</i>	Sep-Feb	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.		Spw.	Spw., Ad.								Ad.
Large mouth nannygai	<i>Lutjanus malabaricus</i>	Oct-Feb (peak Oct-Dec)	Juv.	Juv.	Juv.	Juv.	Juv.		Ad.	Ad.	Ad., Spw							Ad.
Small mouth nannygai	<i>Lutjanus erythropterus</i>	Sep-Apr	Juv.	Juv.	Juv.	Juv.	Juv.		Ad., Juv.	Juv., Ad.	Ad., Spw							
Spanish mackerel	<i>Scomberomorus commerson</i>	Oct-Dec	Juv.	Juv.	Juv.	Juv.	Juv.		Juv.		Ad., Spw							Ad., Spw
Grunter/Javelin fish	<i>Pomadasys kaakan</i> , <i>P. argenteus</i>	Sep-Mar (peak Sep-Nov)	Juv., Ad.	Juv., Ad.	Juv., Ad.	Spw.			Juv., Ad.									Ad.
Fingermark bream	<i>Lutjanis johnii</i>	Sep-Apr	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.			Juv.									Ad.
Mangrove jack	<i>Lutjanus argentimaculatus</i>	Sep-Mar (peak Dec)	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.			Juv., Ad.				Spw.					Ad.
Yellowfin bream	<i>Acanthopagrus australis</i>	Jun-Aug	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.			Spw., Ad.									Ad.
Estuary cod	<i>Epinephelus malabaricus</i>	Mar-Jun	Juv.	Juv.	Juv.	Juv.												Ad.
Queenfish	<i>Seriphus Politus</i>	No data	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.						Spw.?						Ad.



Queenfish	<i>Seriphus Politus</i>	No data	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Ad.
Giant trevally	<i>Caranx ignobilis</i>	Warmer months and linked to the lunar cycle	Juv.	Juv.	Juv.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Ad.	Spw., Ad.
Grey mackerel	<i>Scomberomorus semifasciatus</i>	Sept - Jan		Juv.	Juv.	Juv.	Juv.	Juv.	Juv.	Juv.	Juv.	Ad.	Ad.
Sea mullet	<i>Mugil cephalus</i>	Mar - Jun	Juv. Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Ad., Spw?	Ad.
Hamilton's anchovy	<i>Thryssa hamiltonii</i>	No data		Juv.	Juv.	Juv., Ad.	Juv., Ad.	Juv.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.
Moses perch	<i>Lutjanus russelli</i>	No data	Juv.	Juv.	Juv.	Juv., Ad.	Juv., Ad.	Juv.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Ad.	Ad.
Snapper	<i>Pagrus auratus</i>	May - Aug		Juv.	Juv.	Juv.	Juv.	Juv.	Juv.	Juv.	Juv., Ad.	Ad.	Juv., Ad., Spw
Sweetlip	<i>Lethrinus laticaudis, Lethrinus miniatus</i>	All year	Juv.	Juv.	Juv.	Juv.	Juv.	Juv.	Juv.	Juv.	Juv.	Ad.	Juv., Ad., Spw
Saltwater bony bream	<i>Nematalosa erebi</i>	Sep – May (intensified by flooding)	Juv.	Juv.	Juv.	Juv.	Juv.	Juv.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Ad.	Juv., Ad.
Burnett salmon	<i>Polydactylus sheridani</i>	Oct – Mar (peak Dec)	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.
Silver jewfish	<i>Nibea soldado</i>	No data		Juv.	Juv.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Ad., Spw	Juv., Ad.
Diamond scale mullet	<i>Liza vaigiensis</i>	Dec - Feb	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Ad., Spw	Juv., Ad.
<b>Shellfish</b>	<b>Crustacea</b>												
Mud crab	<i>Scylla serrata</i>	Spring to early Autumn	Juv., Ad.	Juv.	Juv.	Juv.	Juv.	Juv.	Juv.	Juv.	Juv.	Spw.	
Tiger prawn	<i>Penaeus esculentus, P. semisulcatus</i>	All year	Juv.	Juv.	Juv.	Juv.	Juv.	Juv.	Juv.	Juv.	Juv.	Ad.	Ad., Spw
Endeavour prawn	<i>Metapenaeus endeavouri, M. ensis</i>	All year (peak in spring)	Juv.	Juv.	Juv.	Juv.	Juv.	Juv.	Juv.	Juv.	Juv.	Ad., Spw.	
Banana prawn	<i>Fenneropenaeus merguensis</i>	All year (peaks based on the monsoon: late dry season (Sep-Nov) and late wet season (Mar-May))	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Juv., Ad.	Ad., Spw	

Commercial fish catch data (DAFF 2013) for the study area shows that there was great inter-annual variability in commercial fish production in both net (**Figure B7.4.9.1b**) and line (**Figure B7.4.9.1c**) fisheries. All commercial fisheries data presented in this report refer to the Queensland commercial fisheries grid H16, where commercial fishing operations are concentrated in the near shore sub-grids in close proximity to Cairns, as shown in **Appendix D13**. Note that the study area comprises only a portion (<25 percent) of grid H16. Also, this grid includes the regulated waters of Trinity Inlet and Barron River, where the number of commercial licenses has been reduced in recent years and those remaining are not expected to be renewed (contributing to reduced catches in recent years).

For recreational fishing, results from the 2010 state-wide survey (Taylor *et al.* 2012, **Figure B7.4.9.1d**) show coral trout, snapper and pikey bream to be the most commonly caught (and harvested) fish species in Cairns coastal waters. Helmke *et al.* (2003) interrogated the Queensland RFISH database for 1997 and 1999, and found that other key recreational target fish species for Trinity Inlet and the harbour include bream, cod, grunter, mackerel, mangrove jack, sardine, sweetlip and whiting (**Table B7.4.9.1c**).

**Figure B7.4.9.1b Net Fishery Commercial Fish Catch (Tonnes) and Rainfall for the Cairns area (Cell H16)**  
 (Data Source: DAFF Unpublished Data 2013)




Figure B7.4.9.1c Line Fishery Commercial Fish Catch (Tonnes) and Rainfall for the Cairns area (Cell H16) (Data Source: DAFF Unpublished Data 2013, Nil Line Data available after 2005)





Figure B7.4.9.1d Estimated Recreational Fishing Catch for Cairns Coastal Waters in 2010 State-Wide Recreational Fishing Survey (Source: Taylor et al. 2012)



**Table B7.4.9.1c Number of Fish Kept and Numbers Released, for each Species Recorded in the RFISH Database for Trinity Inlet and Cairns Harbour in 1997 and 1999 (from Helmke et al. 2003)**

Common Name	Totals		Kept		Released	
	1997	1999	1997	1999	1997	1999
archerfish	3	0	0	0	3	0
barramundi	7	8	2	5	5	3
batfish	0	6	0	5	0	1
bream	27	93	5	31	22	62
catfish	6	41	1	1	5	40
cod	22	11	10	4	12	7
coral trout	0	8	0	2	0	6
crab	0	330	0	65	0	266
diamond-fish/butterfish	8	0	8	0	0	0
eel	0	3	0	0	0	3
emperors	2	0	0	0	2	0
flathead	4	2	2	1	2	1
grunter	4	117	0	26	4	91
jew	28	5	0	2	28	3
leatherjackets	5	0	0	0	5	0
mackerel	3	55	3	38	0	17
mangrove jack	15	43	1	33	14	10
moses perch	20	4	2	0	18	4
queenfish	1	2	0	0	1	2
remora	0	1	0	0	0	1
sardine	0	30	0	30	0	0
shark	14	19	0	5	14	14
sickle fish	1	10	1	0	0	10
snapper	25	47	2	10	23	37
stingray	1	3	0	0	1	3
sweetlip	31	5	7	3	24	2
threadfin	1	1	1	1	0	0
toad fish	1	2	0	0	1	2
trevally	0	7	0	4	0	3
Triggerfishes	5	0	0	0	5	0
tripod fish	0	1	0	0	0	1
white spotted guitarfish	1	0	0	0	1	0
whiting	0	14	0	11	0	3
<b>Totals</b>	<b>235</b>	<b>868</b>	45	<b>276</b>	<b>190</b>	<b>592</b>


### B7.4.9.2 Peneid Prawn Assemblages

Peneid prawns represented a significant proportion of the commercial fishery in the Cairns area. Tiger, Endeavour and banana prawns represented, respectively, 24, nine and two percent of the total commercial catch by weight, but together accounted for 56 percent of the monetary (gross value of production) value of the commercial fishery. Cairns also represents a key Australian location for the harvest of live prawns which are collected to supply broodstock for the prawn aquaculture industry. Generally, the mosaic of high quality wetland habitats (e.g. extensive juvenile prawn nursery habitat amongst intertidal seagrass meadows) together with the large area of flat, trawlable grounds, are important conditions that together contribute to high prawn productivity. Note that the present poor condition and extent of seagrass meadows are likely associated with the comparatively reduced abundances of juvenile prawns inshore.

Coles et al. (1993) sampled peneid prawn assemblages within the Cairns harbour monthly between 1980 and 1987. Sites were positioned in seagrass meadows and unvegetated sediments, and were sampled using standardised hauls with a beam trawler.

Twenty peneid prawn species were recorded, of which nine are marketed commercially. The most abundant commercial species recorded were tiger prawns *Penaeus esculentus* and *P. semisulcatus*, and the endeavour prawn *Metapenaeus endeavouri*. All collected prawns were juveniles. Significantly higher numbers of commercially important prawns were recorded in seagrass than in unvegetated habitats. Furthermore, higher abundances of juvenile tiger prawns were recorded during summer, which is the key recruitment period for these species (Figure B7.4.9.2a).


**Figure B7.4.9.2a Means and Standard Errors of Numbers of Juvenile Prawns of (a) Endeavour Prawn *Metapenaeus endeavouri*, (b) Brown Tiger Prawn *Penaeus esculentus* and (c) Green Tiger Prawn *P. semisulcatus* caught in each Month in Cairns Harbour 1980-87 (Coles et al. 1993)**



Commercial catch data for the period 1988 to 2011 are presented in Figure B7.4.9.2b. Consistent with patterns observed in juveniles prawns by Coles et al. (1993), tiger prawns (49 percent of the total catch) and Endeavour prawns (34 percent of the catch) dominated commercial prawn catches, followed by banana prawns (12 percent of the catch). This catch pattern was relatively consistent from among years, except in 1992 and 2000 when tiger and Endeavour prawn catches were similar.

Total annual tiger prawn catch tended to be greater in years following higher rainfall, whereas no such association was found in Endeavour prawn catches. Positive (and negative) correlations between commercial fish/shellfish catches and rainfall have been observed by a number of workers (Pen and Caputi 1986; Gillanders and Kingsford 2002; Meager 2003). Catchment rainfall can affect catches in a number of ways including (i) affecting 'catch-ability' of fish/shellfish (ii) increasing nutrient/food availability for recruits; (iii) flushing fish/shellfish from estuarine areas (Loneragan and Bunn 1999).

**Figure B7.4.9.2b Commercial Prawn Catch and Total Commercial Catch (All Species) for the Cairns Area (Cell H16)**  
(Data Source: DAFF Unpublished Data 2013)



### B7.4.9.3 Crabs and Other Shellfish

Mud crabs (*Scylla serrata*) primarily inhabit mangrove forests during their juveniles and adult life-cycle stages. Egg bearing females migrate from inshore mangrove areas to deep offshore waters for spawning (Heasman *et al.*, 1985; Hill, 1994). Spawning of mud crabs during the warmer months, and female mud crabs then return to estuaries after spawning (Hill 1994). Mud crabs would migrate from Trinity Inlet and through the harbour on the way to and from offshore spawning areas.

The numerous mangrove-lined creeks within Trinity Inlet represent locally important mud crab foraging and nursery habitat, as is apparent from commercial mud crab production values shown in **Figure B7.4.9.3a**. There was great variability over time, which was partly due to changes in crabbing effort. While the mangrove areas of Trinity Inlet provide an important habitat role for local mud crabs, crabs are actually widely distributed across the area, with sites at both western (e.g. Ellie Point) and eastern (e.g. Bessie Point) Cairns harbour previously shown to support high mud crab catch rates, relative to Trinity Inlet (Helmke *et al.* 2003, **Figure B7.4.9.3b**).

In terms of other commercially harvested shellfish species, bugs and to a lesser extent squid and scallops represented a locally important component of the fishery within the study area (Figure B7.4.9.3b). There is little life history information available for Far North Queensland populations of these species. Bug species have different habitat requirements, with mud bugs (*Thenus orientalis*<sup>1</sup>) occurring in turbid nearshore waters to a depth of 60 m, over sand, mud and gravel substrata (Kailola *et al.* 1993). Sand bugs (Thenus species) prefer clear deep water (Kailola *et al.* 1993), and are unlikely to present in the study area. Mud bugs are likely to occur in subtidal soft sediment habitats throughout Cairns harbour, and would spawn in offshore waters during spring-summer. Bugs and squid would also occur on soft sediment habitat throughout the harbour.

Mud crabs, in particular, are a key recreational target species, being the most commonly recreationally harvested species in 1999 (**Table B7.4.9.1c**) and fourth commonly harvested species in 2010 (**Figure B7.4.9.3b**), based on the available reported information.

<sup>1</sup> Note that DAFF data lists bugs as 'whole', 'mixed' or 'Balmain', with terms varying between years. It is more likely that the species present in Cairns is Moreton Bay bug, which is abundant in the region.

Figure B7.4.9.3a Commercial Mud Crab Catch Data (Tonnes) and Days of Fishing Effort for the Cairns Area (Cell H16) (Data Source: DAFF Unpublished Data 2013; not available 2006 or 2010 onwards)






Figure B7.4.9.3b Commercial Catch Data (Tonnes) for Other Shellfish in the Cairns area (Cell H16) (Data Source: DAFF Unpublished Data 2013)



**Figure B7.4.9.3b Total and Legal Catch Rate (Number Caught per Pot Lift) of Mud Crabs between March 1997 and March 2000 at Trinity Inlet and Cairns Harbour (from Helmke et al. 2003)**



#### B7.4.9.4 Fisheries Habitat Values of the Study Area

**Table B7.4.9.1b** shows that most key fisheries species found in the study area are not found exclusively in any one habitat type during any part of their life-cycle. Rather, these species have relatively plastic habitat requirements, and are typically found in a variety of habitat types. Banana prawns were the only habitat specialists recorded in the study site, and are typically found in mangrove during their juvenile stages (Staples et al. 1985).

While it is apparent that the habitat types supported in the study area are of importance to fisheries productivity, there are few empirical studies demonstrating that habitat resource availability limits (c.f. influences) estuarine fish and nekto-benthic crustacean population abundances. Most studies tend to suggest that non-equilibrium processes (e.g. predation, mortality) are probably more important in controlling local-scale fish abundances rather than density-dependent habitat or food limitation during the post-settlement phase. It is therefore not especially practical to consider that all patches of a particular habitat type have equal value as a fisheries resource (see for example Tibbetts and Connolly 1998), or indeed that the removal of an area of habitat will result in a commensurate reduction in fisheries productivity.

For marine species with complex life cycles defining the spatial characteristics (i.e. location, shape, surface area or volume) of functional nurseries is a complex task because the locations (and times) of suitable habitats are determined by a range of biological processes, many of which are stochastic in nature. Ultimately functional nursery habitats are those which produce reproductive adults that contribute progeny to the next generation.



Helmke *et al.* (2003) examined the Trinity Inlet and Cairns harbour data in the Suntag database for 1985 to 2000, in order to assess fish movements in the area. It was found that, although most fish remain resident in Trinity Inlet, the inlet also acts as a nursery habitat for species such as Barramundi, which spend the juvenile and sub-adult phases of their life cycle in fresh and estuarine waters (Helmke *et al.* 2003). The few recaptures recorded outside Trinity Inlet, or having moved into the area included: barramundi moving in from North Johnson, Daintree and Barron Rivers. Russell, *et al.* (2003) conducted tagging studies for the population biology of Mangrove Jack *Lutjanus argentimaculatus* and included findings of movements of tagged fish from Trinity Inlet up to 335 kms north to Stapelton Reef near Princess Charlotte Bay, indicating estuary systems such as Trinity Inlet and the Mulgrave-Russell systems are important nursery habitats for such important recreational species.

When considering the fisheries habitat value of an area, it is important to consider the spatial organisation of habitat patches and types (i.e. degree of fragmentation), together with other attributes such as structural complexity and size of habitat patches, and the degree of tidal inundation/flushing. Recent studies have examined the importance of mangroves, seagrasses and saltmarsh as autotrophic nutritional sources for fish in adjacent unvegetated environments (Melville and Connolly 2005, Connolly and Guest 2004, Melville and Connolly 2003). Melville and Connolly (2003) also demonstrated that even on unvegetated soft sediment habitats, organic matter, particularly from seagrasses, was important as the base of food webs for fish species of commercial significance on adjacent unvegetated mudflats in Moreton Bay. Benthic microalgae also contributed a relatively high proportion of the nutrition of the species examined.

Seagrass beds also represent an important (and in some case obligatory) habitat resource for many nektobenthic crustaceans and some fish of commercial significance (see, for example, Bell and Pollard 1989, Connolly, *et al.* 1999, Edgar and Kirkman 1989) (Bell, *et al.* 1988, Haywood, *et al.* 1995). Furthermore, there is an emerging view that fish and nektobenthic crustacean community structure in mangroves and unvegetated habitats is influenced by their proximity to seagrass beds (e.g. Jelbart 2004, Olds 2002). Studies by Olds (2002) in Moreton Bay and Jelbart (2004) in central NSW both found that the suite of species inhabiting seagrass varied with distance from mangroves, whereby seagrass beds (particularly dense beds – Olds 2002) in close proximity to mangroves tend to contain more abundant nekton assemblages than seagrass remote from mangroves.

On this basis, the following broad conclusions are applicable to habitat values of the study area:

- Extensive areas of intertidal habitat are present in the study area, including well-flushed mangals, tidal channels with undercut banks (present along Trinity Inlet), and mud flats in the Cairns harbour. These intertidal environments provide shelter and/or foraging areas for fish and nektobenthos during high tide
- Subtidal habitats, which provide refugia during low tide, occur throughout the project area. This is a potentially important factor determining the 'values' of intertidal habitats. It is thought that the risk of predation is increased where fish need to move large distances between intertidal habitat patches (e.g. mangroves, saltmarsh, tidal flats) and permanent tidal pool habitats during low tide (Crowley and Tibbetts in (Tibbetts and Connolly 1998)
- Seagrass habitats in both intertidal and subtidal waters provide an important foraging and nursery habitat, particular where located in close proximity to the mangrove refugia of Trinity Inlet. However, relative to the other habitats available, seagrass extent and condition can fluctuate greatly over time in response to climate influences
- Limited areas of saltmarsh plants. Saltmarsh communities within the study site are inundated tidally during high water spiring events, and are known from case-studies elsewhere to provide functional habitats and foraging areas for a range of fish (typically small-bodied non-commercial species) and nektobenthic crustaceans (including Penaeid prawns and non-commercial crab species) of indirect and direct fisheries value (e.g. Morton, *et al.* 1987, Mousalli and Connolly 1998).

## B7.5 Assessment of Potential Impacts

### B7.5.1 Assessment Basis

A risk-based approach was adopted in this environmental impact assessment. This is based on the identification of potential impacting processes and characterisation of the likely level of impact to the existing environment. The risk assessment process is described in **Chapter A1, Project Introduction** of this EIS. For the purposes of this Marine Ecology assessment, impacts levels and risks were defined on the basis of the following:

- **Significance of Impact** – made up of assessment of the intensity, scale (geographic extent), duration of impacts and sensitivity of environmental receptors to the impact. Impact significance ratings take into account the conservation management objectives for protected areas (see **Section B7.4.2**) and threatened species (as outlined in the relevant recovery plans for species listed in **Section B7.4.7**). **Table B7.5.1a** is a summary of the categories used to define impact significance

- **Likelihood of Impact** – which assesses the probability of the impact occurring. **Table B7.5.1b** is a summary of the categories used to define impact likelihood
- **Risk rating** – which assesses the level of risk for key impacting processes. The risk rating was generated from the Magnitude and Likelihood scores, based on the matrix presented in **Table B7.5.1c**.

When determining the most appropriate impact significance during the impact assessment process, a number of the terms used in the above significance criteria definitions were further defined by elements of the impact assessment methods used elsewhere in this EIS. Most notably, this includes the ‘zones of impact’ assigned to impacting processes associated with water quality (**Chapter B5, Marine Water Quality**) and sediment deposition (**Chapter B3, Coastal Processes**), which take into account the relevant project-specific ecological threshold values applied for this EIS. In terms of water quality and sediment deposition impacts to ecological receptors, these zones are described as:

- **Zone of Influence** – extent of detectable plume, but no predicted ecological impacts
- **Zone of Low to Moderate Impact** – water quality may be pushed beyond natural variation potentially resulting in sub-lethal impacts to ecological receptors and/or mortality with recovery between six months (lower end of range) to 24 months (upper end of range)
- **Zone of High Impact** – water quality would most likely be pushed beyond natural variation (excluding extreme weather events) potentially resulting in mortality of ecological receptors with recovery greater than 24 months.

Full details on these zones, their determination, and criteria are provided in **Chapter B5, Marine Water Quality**. In brief, threshold values were derived from the most relevant existing information available describing the tolerances of key receptors to environmental modifications (i.e. light) relative to background conditions. Where this ecological assessment is based on modelled outputs (i.e. for impacting processes associated with turbid plumes and sediment deposition), it is recognised that modelling predicts the magnitude and location of effects for varying scenarios. Thus, related ecological effects (in terms of the extent and location) are likewise predictive as opposed to definitive.

Key assumptions and limitations of the impact assessment are outlined in the relevant sections.

**Table B7.5.1a Impact Significance Criteria**

Significance	Definition	How defined
Very High	The impact is considered critical to the decision-making process as it would represent a major change to the ecological character of Cairns harbour and/or the immediate surrounds. This level of impact would be indicated by:	
	<ul style="list-style-type: none"> <li>• Complete, irreversible or otherwise long-term (i.e. greater than decades) loss of all of a given habitat type (presently supported in Cairns harbour or surrounds) which is critical for maintaining high ecological values</li> </ul>	<ul style="list-style-type: none"> <li>• Direct loss (project footprint)</li> </ul>
	<ul style="list-style-type: none"> <li>• Complete, irreversible or otherwise long-term (i.e. greater than decades) loss of any high ecological value community or species (i.e. reef-associated benthos, seagrass, mangroves, high value fisheries species) supported in Cairns harbour or surrounds</li> </ul>	<ul style="list-style-type: none"> <li>• Direct loss (project footprint)</li> </ul>
	<ul style="list-style-type: none"> <li>• Severe impacts to populations of Commonwealth or State listed threatened species, such that their capacity to reproduce and recover is significantly affected.</li> </ul>	<ul style="list-style-type: none"> <li>• Area supports ‘important population’ (as per MNES Guidelines 2013), and action known or likely to cause impacts to population, based on MNES Guidelines (2013).</li> </ul>

Significance	Definition	How defined
High	<p>The impact is considered critical to the</p> <p>The impact is considered important to the decision-making process as it would represent a detectable change to the values that underpin the ecological character of the study area (Cairns harbour and surrounds). This level of impact would be indicated by:</p>	
	<ul style="list-style-type: none"> <li>&gt;10 percent irreversible or medium/long-term (&gt;five years) loss in the extent of any habitat type presently supported in Cairns harbour or surrounds</li> </ul>	<ul style="list-style-type: none"> <li>Direct loss (project footprint)</li> <li>Areas likely to be affected in long-term (&gt;five years) by above ambient deposition levels.</li> </ul>
	<ul style="list-style-type: none"> <li>A detectable significant change to the structure (diversity, richness, composition, etc.) high ecological value community structure (i.e. reef-associated benthos, seagrass, mangroves, high value fisheries species) such that communities lack the capacity to return or recover to a state resembling that prior to being impacted within a timeframe of five years or less</li> </ul>	<ul style="list-style-type: none"> <li>Zone of High Impact – Deposition/turbidity (percentile + ecological thresholds).</li> </ul>
	<ul style="list-style-type: none"> <li>Mortality of a small number of individuals of internationally/nationally threatened species, but no detectable change to population status or the capacity of populations to recover.</li> </ul>	<ul style="list-style-type: none"> <li>Area supports species</li> <li>Loss of individuals from study area (from mortality or permanent abandonment, etc.), but unlikely to result in impacts to population (as per MNES Guidelines).</li> </ul>
Moderate	<p>While important at a state, regional or local scale, these impacts are not likely to be critical decision making issues. This would be indicated by:</p>	
	<ul style="list-style-type: none"> <li>Irreversible reduction in the extent of a key high ecological value habitat type presently supported in Cairns harbour or surrounds.</li> </ul>	<ul style="list-style-type: none"> <li>Direct loss (project footprint map)</li> <li>Areas likely to be affected in long-term (&gt;five years) by above ambient deposition levels.</li> </ul>
	<ul style="list-style-type: none"> <li>A detectable significant change to the structure (diversity, richness, composition, etc.) of a high ecological value community structure (i.e. reef-associated benthos, seagrass, mangroves, high value fisheries species), but recovery to a state resembling that prior to being impacted within a timeframe of five years or less</li> </ul>	<ul style="list-style-type: none"> <li>Zone of Low to Moderate Impact (seagrass, mangroves, corals).</li> </ul>
	<ul style="list-style-type: none"> <li>Temporary loss of life history function for threatened species, or species of high fisheries or otherwise ecological value, but no detectable change in their population status at local (study area and surrounds) spatial scales (e.g. once off interruption of breeding or spawning, not necessarily affecting all of local population).</li> </ul>	<ul style="list-style-type: none"> <li>Area supports species</li> <li>Loss of individuals from study area (from mortality, long-term or temporary abandonment etc.), but unlikely to result in impacts to any local population (as per MNES Guidelines).</li> </ul>

Significance	Definition	How defined
Minor	Impacts are recognisable/detectable but acceptable. These impacts are unlikely to be of importance in the decision making process. Nevertheless, they are relevant in the consideration of standard mitigation measures. This would be indicated by:	<ul style="list-style-type: none"> <li>• Zone of Influence (99th percentile &gt; 1 NTU or &gt; cm/mg/day).</li> </ul>
	<ul style="list-style-type: none"> <li>• Temporary (i.e. within 12 months of completion of dredging) changes to sediment type and soft sediment benthic communities at local scale (measured at scale of 10s to 100s of metres)</li> <li>• Short term (i.e. duration of dredge campaign, less than one year) changes to the distribution of threatened species or species of high fisheries significance (i.e. avoidance of areas), but no long-term effects to local population status.</li> </ul>	
Negligible	Minimal change to the existing situation. This could include, for example, impacts that are below levels of detection, impacts that are within the normal bounds of variation, or impacts that are within the margin of forecasting error.	<ul style="list-style-type: none"> <li>• Zone of Influence</li> </ul>

Table B7.5.1b Likelihood of Impact

Likelihood of Impacts	Risk Probability Categories
Highly Unlikely	Highly unlikely to occur but theoretically possible.
Unlikely	May occur during construction of the project but probability well below 50 percent; unlikely but not negligible.
Possible	Less likely than not but still appreciable; probability of about 50 percent.
Likely	Likely to occur during construction or during a 12 month timeframe; probability greater than 50 percent.
Almost Certain	Very likely to occur as a result of the proposed project construction and/or operations; could occur multiple times during relevant impacting period.

Table B7.5.1c Risk Matrix

Likelihood	Significance				
	Negligible	Minor	Moderate	High	Very High
Highly Unlikely/ Rare	Negligible	Negligible	Low	Medium	High
Unlikely	Negligible	Low	Low	Medium	High
Possible	Negligible	Low	Medium	Medium	High
Likely	Negligible	Medium	Medium	High	Extreme
Almost Certain	Low	Medium	High	Extreme	Extreme

Table B7.5.1d Risk Rating Legend

<b>Extreme Risk</b>	An issue requiring change in project scope; almost certain to result in a 'significant' impact on a Matter of National or State Environmental Significance
<b>High Risk</b>	An issue requiring further detailed investigation and planning to manage and reduce risk; likely to result in a 'significant' impact on a Matter of National or State Environmental Significance
<b>Medium Risk</b>	An issue requiring project specific controls and procedures to manage
<b>Low Risk</b>	Manageable by standard mitigation and similar operating procedures
<b>Negligible Risk</b>	No additional management required

## B7.5.2 Overview

Three major components of the project are considered in this assessment of potential impacts to marine ecology values, namely:

- Widening and deepening of the existing outer shipping channel which will result in some lengthening of the existing channel
- Widening and deepening of the existing inner port channel and cruise shipping swing basin and establishment of a new shipping swing basin to enable future expansion of the HMAS Cairns Navy base
- Structural upgrade of the existing cruise shipping wharves 1-5 to accommodate larger and heavier cruise ships.

The harbour and channel development works will primarily involve capital dredging of approximately 4.4 million m<sup>3</sup> (Mm<sup>3</sup>) of material, and associated placement of this dredged material at a new designated marine (DMPA). Maintenance dredging of harbour and channel areas during the operational phase of the project is also considered.

**Table B7.5.2a** summarises the key processes for each project component that could impact marine ecological values, during either the construction and/or operational phases of the project. Each of these processes and the resultant ecological effects are detailed in the following sections. **Figure B7.5.2a** provides a visual close-up of the direct disturbance footprint in the inner port and wharf upgrade works, while **Figure B7.5.2b** shows the location of the total direct disturbance footprint for the project, relative to the location of key sensitive marine ecology receptors.


**Table B7.5.2a Marine Ecology Impacting Processes during Construction (C) and Operational (O) Phases**

Phase	Impacting Process	Primary Impact	Secondary Effects	Section
C/O	Dredging and dredged material placement	Temporary loss or mobilisation of benthic fauna.	Change in prey availability for marine fauna.	<b>B7.5.3.1, B7.3.5.2</b>
C		Long term change in benthic habitat conditions and benthic fauna.	Change in prey availability for marine fauna.	<b>B7.5.3.1, B7.3.5.2</b>
C/O		Increased suspended solid concentrations and sedimentation.	Loss or degradation of seagrass and corals.	<b>B7.5.4</b>
C/O		Acoustic effects to marine fauna.	Avoidance of area by marine fauna.	<b>B7.5.5.2</b>
C/O		Direct effects of dredge plant on marine megafauna.	Injury or mortality to marine megafauna.	<b>B7.5.5.1</b>
C/O	Wharf infrastructure development and operation	Direct changes to marine habitat.	Change in prey availability for marine fauna.	<b>B7.5.3.3</b>
C/O		Acoustic effects to marine fauna (e.g. physiological damage, masking of biologically important sounds) associated with construction (e.g. piling) and operational vessel noise and vibration.	Adverse marine fauna behavioural responses, temporary avoidance or displacement of affected area.	<b>B7.5.6 B7.5.5.2</b>
C/O	Increased vessel movements	Increase in boat strike (construction vessels and increased cruise ship movements).	Injury or mortality to marine megafauna.	<b>B7.5.5.1</b>
C/O		Increase potential for marine pest introductions.	Out-competition of native species and loss of biodiversity values.	<b>B7.5.7.1</b>
C/O		Increase in vessel wash and disturbance of seabed habitats, flora and benthic fauna.	Change in prey availability for marine fauna.	<b>B7.5.7.2</b>
C/O	Construction plant and operational lighting	Increased light spill into the marine environment.	Disorientation of marine fauna, particularly marine turtles.	<b>B7.5.5.2</b>
C/O	Increased potential for debris and spills to enter the marine environment	Ingestion or entanglement of marine megafauna. Toxicity effects to marine flora and fauna.	Loss of biodiversity values.	<b>B7.5.8</b>

Figure B7.5.2a Capital Dredging Direct Disturbance Footprint for the Inner Port, Wharf Upgrade Works and Nearshore Sections of the Outer Channel



Figure B7.5.2b Direct Disturbance Footprint for the project, Relative to the Location of Key Sensitive Receptors





**Table B7.5.2b** summarises the approximate area and type of marine habitat affected by direct impacting processes at each impact location, as well as other anticipated direct habitat changes.

**Table B7.5.2b Area of Disturbance within each Impact Location**


Location	Phase	Activity	Effect Type	Habitat Type	Area (ha) Affected
<b>Direct Irreversible Losses and Gains at Wharf</b>					
A	C,	Wharf dolphin structures, associated piles for wharf infrastructure.	Loss of soft sediment habitat. Habitat modification – hard substrate habitat associated with wharf upgrade works (84 piles).	Subtidal soft sediments Hard substrate	0.0054 ha 0.0054 ha (gain)
<b>Direct Habitat Disturbance Associated with Dredging Activities</b>					
B	C, O	Dredging and deepening of inner port (previously dredged areas).	Habitat modification - Increase in depth where capital dredging will occur; disturbance by maintenance dredging (at similar frequency as existing maintenance dredging).	Subtidal soft sediments	45.18 ha
C	C, O	Dredging and deepening of inner port in previously un-dredged areas.	Habitat modification - increase in depth where capital dredging will occur; disturbance by maintenance dredging.	Subtidal soft sediments	21.08 ha
D	C, O	Deepening of the existing outer channel (previously dredged areas).	Habitat modification - increase in depth; disturbance by maintenance dredging (at similar frequency as existing maintenance dredging).	Subtidal soft sediments	99.66 ha
E	C, O	Channel widening and increase in length in previously undredged areas.	Habitat modification - increase in depth; ongoing disturbance by maintenance dredging.	Subtidal soft sediments	64.27 ha
F	C, O	Dredged material placement at the DMPA.	Direct habitat modification due to dredged material placement, and ongoing disturbance through maintenance dredging.	Subtidal soft sediments	269 ha
E	C,O	Decommissioning of existing DMPA.	Cessation of dredged material placement activities at the existing DMPA; potential for longer term stabilisation of benthic communities.	Subtidal soft sediments	269 ha

For assessments based on predictive model outputs of dredging scenarios, the following scenarios are considered (in line with **Chapter 3, Coastal Processes** and **Appendix D4, Water Quality Model Development and Calibration Report**):

- Base Case Capital Program
- Alternative Case Capital Program
- Constrained Overflow (Worst Case) A – limited overflow when dredging soft material in channel
- Constrained Overflow (Worst Case) B – limited overflow when encountering stiffer than expected material in channel.

The difference between the 'base case' and the 'alternative case' is the use of different dredging equipment in the inner port, and these cases represent the 'likely best case' modelling scenario (no overflow dredging) while the constrained overflow dredging cases represent the 'likely worst case' modelling scenario (Figure B7.5.2c).

Figure B7.5.2c Conceptual Diagram of Modelling Scenarios



### B7.5.3 Direct Modification of Benthic Habitats and Communities from Dredging, Dredged Material Placement and Wharf Upgrades

This section describes direct impacts to benthic habitats and communities due to dredging activities, dredged material placement and wharf upgrade works. Note that impacts associated with turbid plume generation and flow-on effects to marine ecology are considered in Section B7.5.4.

#### B7.5.3.1 Dredging of the Channel, Inner Harbour and Swing Basins

##### Dredging Specifications

Figure B7.5.2b shows the proposed dredging footprint. For capital works, dredging of the outer channel will widen the existing channel from 90 m to 130 m, deepen the channel from -8.3 m LAT to -9.4 m LAT, and extend the existing 11.2 km channel by approximately one km offshore. Practically, the outer channel will be dredged wider to accommodate channel batters (typically one in four slope), and deeper in some areas to allow siltation between maintenance dredging campaigns (Chapter B3, Coastal Processes).

The inner channel extends for 2.4 km and has variable widths incorporating bends and swing basins. Capital dredging here will expand the existing Crystal Swing Basin for use by cruise ships, relocate the existing main swing basin further south (to be designated Smith's Creek Swing Basin), and increase the width and depth of the existing inner port channel along its full length.

Based on the anticipated dredge volumes and materials, it is anticipated that the dredging program will take approximately 21 (Alternative Case) to 34 (Base Case) weeks, with the most likely dredging program estimated at 23 weeks. Future maintenance dredging requirements are expected to increase from the existing annual volume of 350,000 m<sup>3</sup> to an estimated annual volume of 450,000 m<sup>3</sup> (Chapter B3, Coastal Processes).

##### Benthic Habitat Modification

In total, capital dredging will involve the extraction of approximately 230ha of soft, unconsolidated sediment, of which 145ha is presently subject to annual disturbance due to existing maintenance of the channel and inner port. This will result in the direct removal of soft sediment habitat and biota from within the dredge footprint areas of the existing assessed and approved channel structure. Hence, it is proposed that this project will impact approximately 85ha that has not previously been dredged, namely the extension to the seaward end of the channel and the small batter slope/swing basin extensions within the inner port. No other habitat types are present within the indicative dredge footprint, although it does overlap with the cumulative historic extent of seagrass (refer Section B7.4.4), such that there is the potential that seagrass could re-establish here prior to dredging.

It is expected that dredging to widen and extend the outer channel and inner port area will create benthic habitat conditions that are similar to those found within the existing outer channel and previously dredged areas of the inner port. As discussed in Section B7.4.6, existing benthic habitats and macroinvertebrate assemblages within the outer channel are highly simplified and have low diversity compared to adjacent undredged areas. The existing outer channel and inner port are subject to ongoing disturbance as a result of maintenance dredging.

Water depth will also increase, typically by approximately one – 1.5 m throughout the dredge footprint. The increase in water depth will represent a permanent change in habitat conditions as these depths will be sustained by ongoing maintenance dredging campaigns throughout the operational phase. The following physical habitat responses are predicted:

- Given the increase in water depth, benthic habitats within the dredge footprint are expected to receive slightly lower light levels than present due to light attenuation with depth. This is not considered likely to adversely affect habitat conditions, given that no notable photosynthetic benthos are known to presently occur within the dredge footprint (although seagrass may occur in future if it establishes prior to dredging)
- Localised changes to bed stability on the batter slopes of the dredge footprint (see **Chapter B3, Coastal Processes**)
- Highly localised and minor changes in the speed and direction of currents after development completion. For example, the highest magnitude changes in tidal current flow velocities are not large (generally  $\pm 0.1$  m/s, refer **Chapter B3, Coastal Processes**) but could still lead to minor alterations to local scour or sediment accretion rates at discrete locations (depending on the ultimate dredging scenario and tidal conditions).

### Effects to Benthic Communities

Initially, dredging will cause a temporary loss of biota from within the dredge footprint, since benthic communities typically inhabit the surface sediments that will be extracted by dredging. Biota will soon recolonise the dredge footprint but will continue to be regularly subject to similar disturbance through the ongoing annual maintenance dredging regime.

While in this modified state, it would be expected that benthic communities within both the existing channel/harbour and proposed new dredge areas (i.e. channel widening/extension area and parts of the inner port that have not previously been dredged) will support similar benthic communities and ecological functions as that currently found in the existing channels.

#### ***Benthic Fauna***

As discussed in **Section B7.4.6**, benthic fauna communities within the proposed dredge footprint are largely simplified, with a lower fauna abundance and diversity compared to soft sediment habitats elsewhere in the study area. This relates to both epifauna and infauna communities, and is largely associated with much of the dredge footprint having been exposed to past dredging effects, either directly, or by being located immediately adjacent to previously dredged areas. No reef communities or other features of high fauna biodiversity value occur in the proposed newly dredged areas.

Recolonisation of benthic fauna to a dredged area may occur via several processes including:

- Passive recolonisation, involving the passive settlement of entrained or otherwise resuspended organisms (Morton 1977)
- Larval settlement by planktonic organisms (Skilleter 1998)
- Post-colonisation invasion of the dredged area by adult and juvenile fauna from neighbouring undisturbed areas (noting rates of colonisation dependent on the mobility of the animals present in adjacent areas).

Initial passive recolonisation of dredged areas may occur immediately after dredging, followed shortly by the commencement of recolonisation through larval dispersal or active invasion (within hours to days) (WBM 2004). While commencement of initial recolonisation will occur in a short time frame, 'recovery' (functional recovery in terms of a return to comparable numbers of species and total individuals) for areas that have not previously been dredged would be in the order of months to years but will ultimately be limited by the frequency and timing of maintenance dredging (i.e. maintenance dredging fosters a continuous cycle of disturbance and recovery, such that communities remain in a state of flux). However, such areas are subject to natural disturbances from cyclones and would have significant recovery potential. As such, areas of the dredge footprint that have not previously been dredged can be expected to undergo a long-term shift in community composition, whereby for example, the more robust or opportunistic species would contribute proportionately more to total fauna abundances. On the whole, and throughout the longer term operational phase, benthic fauna communities across the dredge footprint will likely reflect those currently inhabiting the existing dredged areas, taking into account additional external community drivers (i.e. noting that a natural gradient would also occur along the length of the dredge footprint, in line with habitat variables such as sediment grain size and organic content).

## Seagrass

Notable beds of seagrass do not currently occur within the proposed dredge footprint, although past surveys have repeatedly mapped seagrass at one outer harbour section of the dredge footprint (Rasheed *et al.* 2013). When present, the seagrass occurring here was dominated by *Halodule uninervis*, and at times was also comprised of *Cymodocea rotundata*, similar to other seagrass beds previously mapped on the eastern side of the existing channel (Rasheed *et al.* 2013).

While notable seagrass beds do not presently occur, smaller isolated patches of seagrass may occur within the dredge footprint either currently, or from time to time throughout the life of the project, particularly in clearer offshore waters. Based on previous surveys it is assumed that any such patches of seagrass would be sparse and highly variable in their extent and density, in response to both seasonal growth patterns and inter-annual cyclic changes (as described in **Section B7.4.4**). Given the assumed small, sparse and/or ephemeral nature of any seagrass occurring in the dredge footprint, such areas would be considered to be of marginal habitat value to seagrass-dependent marine fauna. Therefore, while direct removal or disturbance of any minor areas of seagrass occurring in the dredge footprint would occur, flow-on effects are not expected.

## Secondary Effects

The change in habitat conditions in the dredge channel is predicted to have highly localised secondary effects to marine flora and fauna. Alterations in the composition and abundance of benthic fauna assemblages can be expected within the dredged area immediately after dredging (i.e. prior to recolonisation), resulting in a temporary loss of prey items for fish and invertebrates. This is expected to result in temporary, localised impacts to fish feeding activities.

Given much of the dredge footprint is already subject to ongoing maintenance dredging, this area does not usually contain large or dense seagrass areas. Hence, any seagrass present is unlikely to provide a critical foraging function for fish, green turtles, dugongs (or other seagrass dependent marine species) compared to the more extensive seagrass meadows normally occurring elsewhere in Cairns harbour (i.e. in the vicinity of Cairns Esplanade and Bessie Point).

Overall, modifications to benthic habitats and communities in the proposed channel extension area are expected to initially result in highly localised reductions in benthic fauna richness and abundance. These communities will begin to recover (possibly commencing immediately after dredging) but will continue to fluctuate in response to maintenance dredging and natural ambient disturbances from extreme weather events, similar to the present situation in maintained dredged areas. Overall, these changes are not expected to cause detectable flow-on effects to other ecosystem components or functions throughout the study area, beyond the dredge footprint.

Note that for marine protected areas, dredging will encroach on the Trinity Inlet FHA and GBR Coast State Marine Park General Use Zone. These effects and corresponding mitigation measures are detailed in **Chapter B2, Nature Conservation Areas**. No dredging will occur within the GBRMP.

### B7.5.3.2 Placement of Dredged Material

It is estimated that the total capital dredge volume will be approximately 4.4 Mm<sup>3</sup>, the bulk of which will come from the outer channel. It is proposed that this material will be placed at a new DMPA, a circular area that is one nm in diameter and located approximately 2 km northeast of the existing DMPA. On completion of capital dredging, the seabed height of the new DMPA area will have been raised by an estimated 1.2 m. The placement of this dredged material will result in direct effects to benthic habitats and communities within the DMPA and adjoining areas, as discussed below. It is acknowledged that the DMPA will be the area subject to the most intense environmental effects resulting from dredged material placement and its designation would recognise that such impacts would occur.

#### Within DMPA

Assuming dredged material is placed across all the DMPA, this activity will result in the complete and permanent burial of soft sediment substrates over the entire 2.69 km<sup>2</sup> DMPA area. Reefs and rubble do not occur within the proposed DMPA or immediate surrounds (i.e. within 15 km). Given the approximate 1.2 m thickness of the placed material from capital dredging, this will result in the smothering of any sessile flora (i.e. algae if present) and fauna (e.g. sea pens, macroinvertebrates; refer **Section B7.4.6**) occurring within the DMPA at the time of capital works. If the depth of placed sediment is less in some areas, it is possible that some mobile burrowing fauna will be able to migrate vertically to the new surficial bed height. Increases in bed heights at the DMPA following each maintenance dredging campaign are anticipated to range from approximately 0.14 m (developed case average) to 0.28 m (developed case maximum), as described in **Appendix D4, Water Quality Model Development and Calibration Report**.

Extensive areas of seagrass are not known to occur at the offshore DMPA (**Section B7.4.4**). Seagrass meadows were not detected during surveys for this project at the existing DMPA, although isolated sparse strands of *Halodule uninervis* has previously been recorded in the vicinity of the existing DMPA (Neil *et al.* 2003). Established seagrass meadows are usually restricted to inshore locations in the vicinity of the inner port and outer harbour, noting that seasonally ephemeral seagrass may occur from time to time in deeper waters near the DMPA. Epibenthic communities at the DMPA and similar offshore areas tend to be dominated by sparse macroalgae and occasional epifauna (e.g. gobies, sea pens, crinoids, refer **Section B7.4.6**).

As the capital and ongoing maintenance dredging campaigns will affect different parts of the DMPA at different times, dredged material placement will create a mosaic of patches with different disturbance histories. During and after the capital and maintenance dredge campaign, benthic organisms will colonise the DMPA through the same three mechanisms described above for the recolonisation of the dredged areas. There will, however, be one notable distinction pertaining to passive recolonisation. It is possible that some fauna will survive dredge entrainment and the subsequent placement process, such that passive recolonisation may be by fauna sourced from a location up to 15 km away from the DMPA (i.e. potential for live fauna to be transported from the dredge area).

While opportunistic species and primary colonisers will commence settlement shortly after placement, other less mobile species will take longer to colonise the DMPA. More mobile surface dwelling fauna such as prawns and shrimps, amphipods, isopods and some worms may move from adjacent undisturbed habitats into the DMPA. Most benthic fauna species have a planktonic stage, and in time would colonise the DMPA through larval settlement.

The recovery timescales will depend on the species or assemblage under consideration. In terms of benthic infauna, most studies done in dynamic coastal environments such as the Port of Townsville DMPA have found relatively rapid recolonisation following dredged material placement (measured in time scales of weeks to months). For example, assessments by Cruz-Motta and Collins (2004) at the Port of Townsville found that assemblages in the DMPA were not different from controls three months after dredging, despite impacts being observed one week after dredging. Sampling at Hay Point found no significant differences in benthos between controls and DMPA sites approximately two years after dredging (Hydrobiology Pty Ltd 2003), but unfortunately sampling intervals were too coarse to determine recovery rates in the short term.

Past studies at the Port of Hay Point examined the impacts of capital dredging (and dredged material placement) on seagrass and epibenthic fauna communities (nine months duration), and their subsequent recovery over a 12-month period (Chartrand *et al.* 2009). It was found (Chartrand *et al.* 2009) that one of the two seagrass species (*Halophila decipiens*) present prior to dredging was recorded in the DMPA nine months after the completion of dredging and bed levelling activities, but had very low biomass compared to baseline conditions. It was argued that *H. decipiens* and other deepwater seagrass such as *H. spinulosa*, have adaptations that allow them to rapidly recover following disturbance, assuming ambient conditions post disturbance return to baseline conditions.

Chartrand *et al.* (2009) also found that the sessile epifauna communities at the DMPA had lower abundance than adjacent 'control' locations during dredging, which they speculated was a result of fauna burial, clogging of feeding and respiratory cilia from sedimentation, or by lack of suitable recruitment habitats. However, they also found that abundances of epifauna at the DMPA were higher than at control sites shortly after the completion of dredging. It was speculated that dredging activities enhanced epifauna abundances, possibly in response to an increase in available nutrients and organic matter. Similar reports of benthic enhancement have been reported in other dredge monitoring studies (e.g. Jones and Candy 1981; Poiner and Kennedy 1984; WBM 2004).

### DMPA Surrounds

As discussed in the **Chapter B3, Coastal Processes** and **Appendix D4, Water Quality Model Development and Calibration Report**, waves will tend to remobilise placed sediments, resulting in the dispersion of sediments outside DMPA. Under normal conditions, only fine sediments will be redistributed, whereas heavier sand fractions may be re-mobilised and dispersed during cyclonic events. The dispersal of sediments from the DMPA may occur over long time-frames, subject to the frequency of high energy wave events.

Recent epibenthic surveys undertaken as part of this EIS (refer **Section B7.4.6**) suggest that the placement of dredged material has not had caused detrimental effects to benthic communities in areas immediately adjoining the existing DMPA. To the contrary, McKenna et al. (2013) reported high and medium density epibenthic cover within and immediately adjoining the southern half of the existing DMPA (relative to subtidal soft sediment habitats in other parts of the study area). These were dominated by soft coral and sea pens in high density areas, and echinoids in medium density areas. Similarly, surveys undertaken by BMT WBM to target the existing DMPA and surrounds (see **Section B7.4.6**) indicated:

- Epifauna abundance at sites located in (Sites 7a, 7b) and adjacent to (Sites 8a, 8b) the existing DMPA was rated medium to high, compared to other sites surveyed in the study area. However, community composition differed, with epifauna abundances being comprised of a much higher proportion of sea pens at all four sites
- For infauna communities at these sites disturbed by past dredged material placement activities, both fauna diversity and abundance was moderate, typically on average for all sites surveyed.

These results are consistent with a past study undertaken by Neil *et al.* (2003) who reported that despite a detectable difference in the benthic communities of the existing DMPA and placement affected (northern) surrounds as compared to reference sites, assemblage differences were minor and did not alter the complete assemblage structure of these soft sediment environments. Rather, any differences were only associated with the presence of patchily distributed taxa that occurred in low numbers. Likewise, Worley Parsons (2009) assessed infauna assemblages at the existing DMPA and found minimal impacts attributable to dredged material placement at locations adjacent to the DMPA (i.e. subtle assemblage differences, based on multivariate analysis of community structure). They also confirmed that any impacts were limited to within tens of metres from the DMPA boundary. Similar benthic resilience has been described during longer term studies of DMPAs elsewhere. Monitoring of benthic communities within and adjacent to the Townsville DMPA (Cruz-Motta 2000) indicated that benthic macroinvertebrate communities are resilient to changes in morphodynamics, and that despite a long history of dredged material placement activities, long-term changes in community structure within or adjacent to the DMPA were not observed.

It is therefore expected that the placement of dredged material at the DMPA will not result in major changes to benthic communities outside the DMPA. Monitoring of benthic communities will be undertaken to quantify any changes to communities in the vicinity of the DMPA.

### **B7.5.3.3 Wharf Upgrade Works**

Marine aspects of the wharf upgrade works focus on the installation of 21 independent dolphin structures between existing bents, together with a new fender system every five bents. Each dolphin has four steel piles (totalling 84 piles during construction), concrete pile caps and mooring bollards. Piles are 900 mm in diameter, subject to detailed design, equalling a total base area of 53.76 m<sup>2</sup> for all piles combined.

This area (53.76 m<sup>2</sup>) represents the area of marine habitat and associated benthic infauna communities that will be permanently displaced through the construction of the wharf upgrade works. All of this habitat will be subtidal soft sediment (mud) and represents a small proportion of available soft sediment habitat within the inner port local area (refer sediment class distribution map, **Section B7.4.6**). This area is already in a modified condition due to existing development. As such, this loss is not expected to result in detectable flow-on effects to other local fauna components (e.g. fish, large invertebrates) that rely on benthic infauna as a food resource. The loss of benthic habitat and associated assemblages within the wharf upgrade footprint is irreversible and therefore rated as a moderate impact.

The piles will provide additional artificial hard substrata (i.e. 84 piles in approximately eight m water), which will gradually be colonised by sessile and encrusting biota over time (e.g. algae, attached bivalves, molluscs, bryozoans, etc.), resulting in a benefit in terms of habitat availability for these hard substrate associated communities. It is likely that species richness and biomass of benthic assemblages on the piles will be far greater than on the soft substrate that it replaced. While the piles will act as a fish aggregation device, they are unlikely to increase fisheries productivity except at localised spatial scales.

### **B7.5.3.4 Habitat Changes due to Altered Hydrodynamics**

As discussed in **Chapter B3, Coastal Processes** and **Appendix D4, Water Quality Model Development and Calibration Report**, the project is predicted to result in minor, highly localised changes to hydrodynamics. Within and immediately adjacent to the dredging footprint, depth-averaged current speeds are predicted to increase/decrease (depending on location) within a range of  $\pm 0.2$  m/s. Smaller magnitude reductions in peak current speeds (<10 percent) are predicted more broadly within Trinity Bay.

These minor changes in hydrodynamics would be expected to result in negligible changes to marine habitats and biota.

### B7.5.3.5 Changes to Habitat and Prey Resources for Species of Economic Significance

Two critical considerations when considering the potential impacts of loss or disturbances to benthic assemblages on the foraging of fishery species are: (i) the spatial scale of the impact relative to the total area of habitat available, and (ii) the degree of foraging specialisation exhibited by key fishery species. With respect to loss or changes in prey resource availability, the level of impact will depend on whether the animal has a highly specialised diet, and whether the area affected contains critical food resources.

The total area of soft sediment habitat loss proposed (as described above for dredging, dredged material placement and piling) is relatively small in comparison to the total available soft sediment habitat resource in the harbour, Trinity Inlet and the wider study area. Based on habitat assessments and benthic macroinvertebrate community surveys, none of the potentially affected areas are known to support unique benthic macroinvertebrate or benthic habitats, nor are benthic macroinvertebrate communities within these areas considered to be particularly diverse or abundant compared to adjacent areas (consistent with much of the project footprint already having been subject to past/present disturbance).

Fish species occurring in unvegetated soft sediment habitats are generally recognised as being opportunistic benthic foragers (e.g. Hobday *et al.* 1999). This is demonstrated by how rapidly some fish species occurring in these habitats learn to consume introduced invertebrate species such as bivalves and polychaetes (Hobday *et al.* 1999). Similarly, prawns and crabs of economic significance also have a plastic diet (Dall 1992; Wassenberg and Hill, 1987).

Key commercial and recreational fisheries species potentially occurring at and adjacent to areas of disturbance can be broadly classified as broadly opportunistic species which feed on a wide variety of benthic invertebrates and pelagic fish (Table B7.5.3.5a).

**Table B7.5.3.5a Prey of key harvested species that may overlap spatially with the area of habitat impacts for the project**

Species	Prey	Source
Eastern king prawn, Tiger prawn, Banana prawn	Benthic invertebrates – crustaceans and polychaetes.	Moriarty (1977)
Blue swimmer crab	Benthic invertebrates - crustaceans, molluscs, echinoderms, polychaetes.	Williams (1982), Wassenberg and Hill (1987)
Mud crab	Benthic invertebrates – molluscs, crustaceans, sedentary or moribund fish.	Williams (1997)
Barramundi	Fish and macrocrustaceans (prawns, crabs, etc.).	Davis (1987)
Threadfin salmon	Demersal and pelagic fish (e.g. ponyfish, flathead, scats, sardines) and macrocrustaceans.	Kailola <i>et al.</i> (1993)
Queenfish	A variety of pelagic fish species and cephalopods.	Kailola <i>et al.</i> (1993)
Sand whiting	Benthic invertebrates – crustaceans, molluscs, polychaetes.	McKay (1992)

Given the opportunistic behaviour of the benthic foragers listed above, together with the small proportion of habitat lost, it is not expected that permanent loss or modification of habitat would lead to a long-term reduction in populations of species of economic significance. However, it would be expected that demersal fish, crabs and prawns will avoid areas that have depauperate benthic macroinvertebrate assemblages as a result of dredging and dredged material placement. This is expected to result in a redistribution of fauna, with animals foraging in other parts of the study area (e.g. adjacent to project footprint) until such times as benthic communities recolonise the disturbed area (i.e. recolonisation will commence immediately in dredged areas and DMPA, but invertebrates communities in such areas will likely remain in a cyclical state of flux in response to ongoing maintenance disturbance).

Mud crabs, blue swimmer crabs and demersal fish utilise a range of soft sediment habitat types. There is no empirical evidence to suggest that these species have a strong association with a particular sediment type. Some correlative preferences for sediment type (i.e. grain size distributions) have been shown for commercial prawn species (e.g. Somers 1994). However, Somers (1994) suggested that variables other than sediment grain size may be more important, particularly factors such as the availability and extent of food and nursery habitats (e.g. seagrass, mangrove and benthic faunal communities). In the context of this project, this means that longer term changes in habitat conditions (e.g. sediment types, water depths) as a result of dredging and dredged material placement activities, and associated changes to benthic macroinvertebrate communities, are not expected to lead to significant impacts to these species of economic significance.

The residual impact is considered to be minor for species of economic significance. Further discussion on fisheries impacts specifically relating to the generation of turbid plumes is provided in **Section B7.5.4.1**.

### **B7.5.3.6 Mitigation and Residual Effects**

Habitat removal and physical habitat alteration within the project footprint is an inherent impact of any project that incorporates a marine dredging component. For the project, considerable effort during the design and EIS phases went to the identification and selection of both a dredge footprint and DMPA site that would minimise new direct ecological effects to marine environmental values. The dredge footprint largely aligns with that of the existing outer channel (albeit wider) and other areas that are already regularly dredged under existing conditions. This reduces the extent of dredging required in greenfield areas. A detailed selection process was also undertaken (including on-land placement options) that incorporated ecological considerations, in order to identify a preferred DMPA site (**Appendix D4, Water Quality Model Development and Calibration Report**).

The Dredge Management Plan (**Chapter C2, Dredge Management Plan**) provides guidance on (i) the mitigation measures that will be adopted to minimise direct impacts to marine flora and fauna; and (ii) monitoring that will be undertaken to validate impact predictions outlined in the EIS. This includes the following relevant strategies and components:

- A bathymetry survey of the DMPA and immediate surrounds will be undertaken immediately prior to capital dredging in order to optimise the dredged material relocation strategy and to describe baseline conditions
- Dredge tracks will be reviewed on an ongoing basis in order to spread the dredged material as evenly as practicable, thereby minimising the potential for high spots to occur
- A bathymetry survey of the DMPA and surrounds will be undertaken to confirm final depths at the completion of the capital dredging campaigns
- A monitoring program will be developed and implemented to assess the effects of dredged material placement and subsequent recovery of benthic communities within and adjacent to the DMPA.

For marine protected areas, note that offsets are proposed to compensate for dredging activities encroaching on these areas (e.g. Trinity Inlet FHA), as detailed in full in **Chapter B2, Nature Conservation Areas**.

The location and extent of physical habitat alteration and disturbance during construction phase dredging is unlikely to affect marine flora or fauna of conservation significance. Still, these effects are considered to represent a **Minor to Moderate impact**, given that these habitat alterations will persist in the long term.

## **B7.5.4 Increased Suspended Sediment Concentrations and Sedimentation from Dredging and Dredged Material Placement**

Both dredging and dredged material placement will generate turbid plumes that will extend over marine areas outside the project footprint. The two key effects of this for consideration in terms of potential impacts to marine ecology relate to:

- Water quality degradation from suspended sediment concentrations increasing turbidity
- Increased sediment deposition from suspended sediments settling out of the water column.

These two items are discussed separately below, based primarily on predicted impacts described in the **Chapter B5, Marine Water Quality** and **Chapter B3, Coastal Processes** for the various dredging scenarios.



### B7.5.4.1 Increased Suspended Sediment Concentrations and Turbidity (Capital Works)

#### Predicted Turbidity

Turbid plumes generated by dredging will reduce light levels on the seabed, which could affect photosynthetic benthic species requiring light for energy production (e.g. seagrass, algae, and soft coral). The actual impact of turbid plumes on these benthic primary producers will depend on whether critical light requirements are met, and consideration of the magnitude, frequency and duration of low light events.

Numerical modelling of turbid plumes has been carried out for a range of scenarios; refer to **Chapter B5, Marine Water Quality** and **Appendix D4, Water Quality Model Development and Calibration Report**. Ecological impact assessments in this section are primarily based on the following key modelling outputs:

- **Base Case Scenario** – Impact of dredging on the 50<sup>th</sup> and 95<sup>th</sup> percentile turbidity (**Figure B7.5.4.1b**, **Figure B7.5.4.1c**) and the corresponding impact zone map (**Figure B7.5.4.1d**)
- **Worst Case Scenario 1** – Impact of dredging on the 50<sup>th</sup> and 95<sup>th</sup> percentile turbidity (**Figure B7.5.4.1e**, **Figure B7.5.4.1f**)
- **Worst Case Scenario 2** – Impact of dredging on the 50<sup>th</sup> and 95<sup>th</sup> percentile turbidity (**Figure B7.5.4.1g**, **Figure B7.5.4.1h**), and the corresponding impact zone map incorporating both worst case one and two scenarios (**Figure B7.5.4.1i**).

Note that the base case scenario assumes no overflow from the TSHD; while the worst case scenarios assume varying levels of overflow from the TSHD, depending on the type of bed materials encountered and during the worst 30 day period (i.e. taking into consideration both climatic and operational factors). In the context of these outputs, impacts to the 95<sup>th</sup> percentile turbidity represent the predicted acute water quality effects above background levels over a short-term period (36 hours); while impacts to the 50<sup>th</sup> percentile turbidity indicate more chronic cumulative water quality effects over a longer (15 day) duration (refer to **Appendix D4, Water Quality Model Development and Calibration Report** for further details on the assessment approach). For comparative purposes, modelling results for an additional alternative case scenario can also be found in **Chapter B5, Marine Water Quality**. The extent, location and magnitude of turbidity plumes will ultimately depend on where a dredge vessel is operating at any given time, what type of dredger is operating, and the meteorological/sea conditions at the time of dredging and placement activities.

Based on the modelled base case scenarios presented here (**Figure B7.5.4.1b**, **Figure B7.5.4.1c**), it is predicted that median turbidity would increase slightly (in the order of two to four NTU) along the northern coastline from near the mouth of Barron River to Yorkeys Knob. The greatest increase to median turbidity would be near the channel dredging area and at the proposed DMPA, which would increase by approximately six NTU. For the acute (95<sup>th</sup> percentile) base case conditions, turbidity is predicted to temporarily increase by approximately 10-20 NTU above background conditions in close proximity to the channel dredging area and the DMPA.

For context, a percentile contour plot showing modelled ambient turbidity (without dredging) during 95<sup>th</sup> percentile conditions is provided in **Figure B7.5.4.1a**. Further modelled ambient turbidity plots are included in **Appendix D4, Water Quality Model Development and Calibration Report**.

In terms of impact zones, the zone of low to moderate impact extends out from the DMPA approximately 2.5 km in a north-west and south-east direction, while the zone of high impact is restricted to within the DMPA (**Chapter B5, Marine Water Quality**). While there are no predicted areas of low to moderate or high impact near the dredging areas, as discussed below there may be some edge effects adjacent to the channel which are not indicated by the modelling results. As such, this assessment assumes that there would be comparable (i.e. a zone of low to moderate impact) edge effects directly adjoining the outside edge of the dredge footprint.

Note that the zones of impact are larger in the offshore area due to less turbid ambient conditions (i.e. plumes would be easier to detect above background).

For the worst case scenarios (**Figure B7.5.4.1e** to **Figure B7.5.4.1h**), dredge plumes would extend from the dredging area in a north-westerly direction along the coastline. Median (50<sup>th</sup> percentile) turbidity would increase slightly (up to two NTU) along the northern coastline up to Taylor's Point (south of Double Island), with the greatest increase near the channel dredging area (up to approximately 14 NTU above background). Under 95<sup>th</sup> percentile conditions, turbidity would increase by approximately 10-30 NTU above background conditions (**Chapter B5, Marine Water Quality**).

The main difference in the zones of impact between the base case scenario and worst case scenario (**Figure B7.5.4.1i**) is the extent of the zone of influence (i.e. extent of detectable plumes) near the coastline. By definition, major impacts to marine flora and fauna are not expected in the zone of influence. There is little difference in the extent of the zone of low to moderate impact and the zone of high impact between the base case and worst case scenarios. These zones are still restricted to the vicinity of the proposed DMPA for the worst case scenario.

Further details of the predicted turbidity changes at specific locations are provided in **Chapter B5, Water Quality**, as are similar results for the other scenarios modelled (e.g. alternative case).

**Figure B7.5.4.1a 95<sup>th</sup> Percentile of Modelled Ambient Turbidity (without dredging)**




Figure B7.5.4.1b Impact of Dredging on 50<sup>th</sup> Percentile Base Case Turbidity (Scale: 2 to 40 NTU)




Figure B7.5.4.1c Impact of Dredging on 95<sup>th</sup> Percentile Base Case Turbidity (Scale: 10 to 200 NTU)




Figure B7.5.4.1d Zone of Impact Map for Base Case Scenario




Figure B7.5.4.1e Worst Case Scenario 1 (Soft Material) - Impact of Dredging on 50<sup>th</sup> Percentile Turbidity (Scale: 2 to 40 NTU)




Figure B7.5.4.1f Worst Case Scenario 1 (Soft Material) - Impact of Dredging on 95th Percentile Turbidity (Scale: 10 to 200 NTU)




Figure B7.5.4.1g Worst Case Scenario 2 (stiff material) - Impact of Dredging on 50th Percentile Turbidity (Scale: 2 to 40 NTU)






Figure B7.5.4.1h Worst Case Scenario 2 (stiff material) - Impact of Dredging on 95th Percentile Turbidity (Scale: 2 to 40 NTU)





Figure B7.5.4.1i Zone of Impact Map for Worst Case Scenario



## Biotic Effects

Existing outer channel and harbour areas are routinely dredged for navigation maintenance purposes. Past studies (Carter *et al.* 2002, Environment North 2005, Worley Parsons 2010) suggest that previous utilisation of the existing DMPA for dredged material placement has not resulted in long-term impacts to nearby sensitive receptors. Biological effects are re-examined below in the context of the project.

Seagrasses and hard corals, as well as other photosynthetic biota (e.g. algae, some soft coral), are considered the key sensitive receptors most vulnerable to turbid plume effects. The zones of low to moderate or high impact are predominantly located over offshore soft sediment habitats in and around the DMPA, and do not coincide with any substantial seagrass meadows, coral reefs or any known high density benthic fauna communities for any of the dredging scenarios. The benthic communities found in these zones could be affected by the higher suspended sediment concentrations, as discussed in **Section B7.5.4.2**. While not expressed by the modelling results, this assessment assumes that there would also be comparable (i.e. a zone of low to moderate impact) edge effects directly adjoining the outside edge of the dredge footprint.

The zone of influence (under any modelled dredge scenario) coincides with known (as mapped in 2013) seagrass meadows and coral reefs (e.g. Cairns Harbour, Double Island). By definition, the zone of influence includes areas where detectable turbidity changes could occur, but major ecological effects are not expected based on known tolerances of sensitive receptors. There are some uncertainties regarding the sensitivities of 'new growth' seagrass (i.e. seagrass that has not been mapped in 2013 but new shoots occur prior to the dredging campaign) and seagrass that is at or near the limits of its tolerance range, and implications are discussed further below.

### Seagrass

Seagrass communities in the vicinity of Cairns harbour are usually dominated by *Halodule uninervis*, *Zostera muelleri*, and to a lesser extent *Halophila ovalis*. In the context of the tolerances of these species to increased turbidity and light attenuation, the following is noted:

- *Zostera muelleri* can survive up to a month at low light levels (five percent surface irradiance) but requires 30 percent surface irradiance for long-term survival, as shown in studies undertaken further south at Cleveland Bay (Grice *et al.* 1996). Studies of seagrasses in tropical regions have shown that *Zostera* spp. have significantly greater light requirements (Grice *et al.* 1996, Bach *et al.* 1998, Collier *et al.* 2009) than other species occurring in the study area such as *Halodule uninervis* and *Halophila* spp. (Freeman *et al.* 2008)
- *Halodule* spp. appear to be reasonably tolerant to light deprivation, with *Halodule pinifolia* surviving up to three to four months following complete light attenuation (Longstaff *et al.* 1999). Note, however, that similar data are not available for the *Halodule uninervis* species occurring in Cairns harbour
- *Halophila ovalis* is among the most sensitive species to light attenuation (Longstaff *et al.* 1999). This species can show signs of stress after several days of complete light attenuation and mortality within 30 days of complete attenuation (Longstaff *et al.* 1999)
- Some seagrass species are able to tolerate episodic pulses of high turbidity over an extended period. For example, Chaterand *et al.* (2012) conducted shading experiments to determine the effects of short pulses of low light (shading) conditions over eight, 12 and 16 week periods on the seagrass *Zostera muelleri*. Significant declines in seagrass were recorded under light deprivation between three and four weeks, however, cyclic shading (two weeks of shade followed by two weeks of light) resulted in no declines after eight weeks. No such studies have been conducted on *Halodule uninervis* or *Halophila ovalis* to date.
- In consultation with JCU, seagrass tolerance values which may be relevant for Cairns were developed to test water quality thresholds in **Chapter B5, Marine Water Quality**. These tolerance values included the following:
  - Total loss of seagrass (zone of high impact) would likely occur if the light requirement (LR) was not met for more than six weeks for *Zostera* (LR = 4.5-12 mol/m<sup>2</sup>/day rolling two week average) and more than 21 days for *Halophila ovalis* (LR = 2.8-4.4 mol/m<sup>2</sup>/day).
  - Declines in seagrass but some recovery (zone of low to moderate impact) would likely occur if the LR was not met for one week (low impact) to six weeks (moderate impact) for *Zostera* and one week (low impact) to three weeks (moderate impact) for *Halophila ovalis* during the growing season (July-Dec).
  - No predicted impacts to seagrass (zone of influence) if light does not fall below LR for *Halophila ovalis* and *Zostera* for more than seven consecutive days.

- In the context of the above, it is important to note that there is little information on the tolerances of seagrass during periods of recovery. While it is thought that new seagrass regrowth (e.g. new shoots, seedlings) would be less resilient to reduced light levels, there is a high degree of uncertainty as to what appropriate thresholds would be. In general, (i) new seedlings/shoots have a low energy store so are more dependent on photosynthesis and would be less resilient to periods of low light; and (ii) new seedlings and shoots would have high energy requirements in order to sustain the high rate of growth required to become established (Jarvis *et al.* 2014; pers. comm. M. Rasheed, 2014)
- As described in **Section B7.4.4**, seagrass meadows in near shore waters of the Great Barrier Reef region typically display a seasonal growth cycle in intertidal and shallow subtidal seagrass meadows (Waycott *et al.* 2005; Unsworth *et al.* 2009), with greatest seagrass growth typically occurring in late spring-summer. In the Cairns region, seagrass (*Zostera*) has been observed to have maximum above-ground biomass in late Spring, maximum leaf canopy height from October to February, and flowering in September and October (McKenzie 1994). Considering all of the most common seagrass species in Cairns, generally the nominal seagrass growth period is thought to be from July to November, peaking from September to October. These trends presumably reflect, at least in part, that seagrass light requirements can vary seasonally (Coles *et al.* 2004), whereby increased growth is a response to optimal growth conditions (light availability).

As discussed above, the zones of low to moderate (major changes to seagrass biomass), high (complete seagrass mortality) impact, and assumed edge effects do not coincide with any known seagrass meadows currently present, although it is possible that occasional ephemeral seagrass plants could occur in this area. Effects would occur if (historical) seagrass meadows in close proximity to the channel and/or predicted impact zones were to recover prior to the commencement of dredging.

Existing seagrass meadows (i.e. seagrass meadows mapped within the study area as of 2013) are not expected to be significantly impacted (i.e. mortality, major loss of biomass) within the zone of influence. Notwithstanding this, as stated above, the impact thresholds adopted are based on tolerance data for 'established' seagrasses elsewhere in tropical Queensland (as well as ambient water quality monitoring data - see **Chapter B5, Marine Water Quality**). As discussed in **Section B7.4.4**, seagrasses in Cairns harbour are in a recovery phase (and/or may be at the time of dredging) following major declines over the last few years. As such, this assessment conservatively assumes that even minor turbidity increases could potentially affect new seagrass growth in recovering areas, particularly in areas adjacent to the channel where turbidity generated by dredging will be greatest. On this basis, potential impacts to recovering seagrass areas are considered to be of moderate significance.

Overall, given (i) the minor to moderate impacts potentially occurring; (ii) the current condition and extent of seagrasses; and (iii) the temporary nature of turbid plumes, it is not believed that the water quality effects resulting from the project will preclude the longer term recovery of seagrass (following large scale declines over the last few years in response to natural disturbance) at the broader Cairns harbour level. Nonetheless, seagrass monitoring will be critical to mitigating any such effects and ensuring that no significant impacts will occur (as listed under mitigation in **Section B7.5.4.4**). Seagrass surveys will be undertaken before dredging works, in order to define where areas of active seagrass recovery (i.e. new shoots, seedlings) are located at the time of dredging works. Further, ongoing monitoring of seagrass condition at both established meadows and recovering areas will form a key component of the reactive monitoring program that will be undertaken during dredging.

In the (unlikely but possible) event that seagrass mortality occurs as a result of increased turbidity, recovery could occur through a number of mechanisms. Seagrass species found in the study area such as *Halodule uninervis* and *Halophila ovalis* have adaptations that can allow relatively rapid growth and recovery following disturbance (Duarte *et al.* 1997). Most notably, they can reproduce both sexually and asexually. Where some shoots or roots remain intact, vegetative growth through their rhizomes can allow significant recovery in a relatively short time frame (Coles *et al.* 2004). For small experimentally manipulated plots of *Zostera* at Cairns, Rasheed, (1999) demonstrated that recolonisation primarily occurred through this vegetative growth from surrounding rhizomes (with 12 month recovery period for small scale – 0.25 m<sup>2</sup> – disturbances). Where mortality is more extensive, recovery would be dependent on germination of seeds stored in the seed bank at that location, or on colonisation by propagules dispersed from elsewhere. Seeds may be able to survive in a dormant condition for two to three years and still remain viable (Campbell and MacKenzie 2004, Orth *et al.* 2006). Recent studies confirm the existence of viable seeds in the Cairns area and the presence of a functioning sediment seed bank (Jarvis *et al.* 2013). However, seed density and viability was considered to be low and it is not known how long this seed bank would be able to persist and remain viable (Jarvis *et al.* 2013) as this is the first time such surveys have been undertaken and the range of natural variability for the local species is yet to be fully understood. Overall, the rate of recovery would be dependent on factors such as the location, magnitude and extent of disturbance, as well as the time of year and environmental conditions during the recovery period.

### *Fish and Nekto-benthic Invertebrates of Commercial Significance*

Most fish expected to occur in the study area have a lateral line system, which assists fish to feed in highly turbid waters. Disturbance of the seafloor by dredging and subsequent dredged material placement will result in mobilisation and entrainment of invertebrates in the water column. This increase in the availability of food resources is expected to lead to an increase in the abundance of fish that feed on invertebrates to the dredging and dredged material placement sites. The increase in small fish could have a localised cascading effect, with piscivorous fish and dolphins also attracted to the dredge and dredged material placement sites. This could therefore result in localised changes to fish distribution and abundance, and potentially higher rates of predation.

Turbid plumes may also result in physiological effects to fish. Jenkins and McKinnon (2006) suggested that very high suspended solid concentrations (e.g. 4000 mg/L) could cause gill blockage and eventually mortality to fish. There are very few documented cases of fish kills resulting solely from turbid plumes, and in any case, such concentrations would only be expected occur only rarely and at highly localised spatial scale (within the immediate vicinity of the dredger). Blaber and Blaber (1980) also suggested that turbidity gradients may aid fish larvae in locating estuarine nursery grounds. Although empirical data are lacking, it is possible that the creation of a turbidity gradient during the recruitment period of key species may lead to larvae being attracted to a region where settlement and recruitment rates are normally low due to lack of suitable estuarine habitat.

As discussed in **Section B7.4.9**, prawns and portunid (mud and sand) crabs represent key species of commercial significance, and utilise both near shore and offshore waters (including parts of the study area) for parts of their life cycle. These species primarily inhabit turbid water environments, and are tolerant of a wide range of turbidity conditions. Direct impacts to prawns as a result of high suspended sediment concentrations and sedimentation are therefore not expected and considered to be a negligible impact.

Impacts to key recreational target species (such as mackerel, grunter etc.) are not predicted to be significant, with some localised short term-impacts predicted, dependant on the timing of dredging along the length of the channel relative to time of year and movement of schools within Trinity Bay.

Further, given photosynthetic epibiota (including seagrass) are sparse within the predicted zones of impact, the epibiota in these areas are not considered to represent a key food resource for species of fisheries significance. Therefore, flow-on effects to fisheries as a result of turbid plumes impacting fish food resources are not expected.

### *Marine Megafauna*

Dolphins are the most commonly occurring cetacean species occurring in the study area (refer **Section B7.4.7.4**) and are capable of successfully foraging in turbid waters. Dolphins often stir up bed sediments when foraging for benthic prey, resulting in limited to no visibility for prey detection. It is thought that dolphins detect prey using echolocation rather than visual cues (Mustoe 2006, 2008). Dugongs have poorly developed eyesight and rely on bristles on their upper lip, rather than visual cues, to detect seagrass food resources. Therefore, high suspended solid concentrations generated by dredging and dredged material placement are not expected to adversely affect foraging success for cetaceans or dugongs.

Sea turtles generally have good eyesight and rely on visual and olfactory cues to detect prey and other food resources (Swimmer et al. 2005). Flatback turtles are known to feed in turbid shallow waters (Robins 1995) and may not be directly affected by turbid plumes generated by dredging and placement. Other species such as green and hawksbill turtle, which feed on seagrass and/or in reef environments, may avoid areas affected by turbid plumes. It is noted, however, that key foraging habitats for these species (i.e. reefs, notable seagrass beds) generally do not coincide with the predicted extent of turbidity impact zones.

### *Other Receptors*

The predicted plumes classified as potentially resulting in ecological effects (i.e. zones of low to moderate and high impacts) do not coincide with the known locations of other sensitive biotic receptors such as coral reef communities. Similarly, the modelled dredge plumes of a level that could cause ecological impacts are not predicted to extend to other habitat types present in the broader study area (i.e. occur in the vicinity of soft sediments or pelagic waters only).

Given the extent and location of potentially impacting plumes within the zones of low to moderate and high impacts, together with the sparseness of photosynthetic epibiota in these offshore areas, detectable flow-on effects to other fauna communities are not anticipated. The sparseness of sensitive receptors (e.g. seagrass, soft corals) suggests they would not be of critical importance as a food source to species of high conservation or fisheries value, and that flow-on effects to additional receptors will be negligible.

Significant impacts to marine protected areas are not predicted, with the exception of minor impacts in the vicinity of the DMPA and adjacent surrounds (GBRMP General Use Zone); refer to **Chapter B2, Nature Conservation Areas** for further discussion.

## B7.5.4.2 Increased Sedimentation (Capital Works)

### Predicted Deposition

Details on the predicted sediment deposition impacts throughout the study area are provided in full in **Chapter B3, Coastal Processes** and **Appendix D4, Water Quality Model Development and Calibration Report**. In brief, sedimentation resulting from the deposition of suspended solids from dredging plumes at both the dredge footprint and DMPA are predicted to occur as shown in **Figure B7.5.4.2a** for the predicted cumulative sediment disposition over the duration of the capital dredging base case; and **Figure B7.5.4.2b** for the cumulative sediment deposition occurring during 12 months of re-suspension after the completion of capital dredging (base case) works.

Over the course of the dredging period (base case), moderate sediment deposition in the order of 1000 mg/cm<sup>2</sup> (equivalent to approximately 10 mm depth) above background sedimentation levels is restricted to areas located in or immediately adjacent to the dredge footprint or DMPA, extending out to a maximum distance of approximately two km to the northwest of the DMPA. Sediment deposition reduces with increasing distance from these areas to a low level cumulative total of 50 mg/cm<sup>2</sup> (approximately 0.5 mm depth, which could not be measured) over the dredge period, which occurs: i) up to 10 km to the northwest of the DMPA, and ii) approximately three km upstream of the dredge footprint in Trinity Inlet. Note that ongoing (post capital dredging) sediment deposition resulting from 12 months of re-suspension is minimal (**Figure B7.5.4.2b**), confined to a maximum of approximately 200 mg/cm<sup>2</sup> (i.e. 2 mm) immediately adjoining to the dredge footprint and less than 50 mg/cm<sup>2</sup> (<0.5 mm) in the DMPA.

To consider these sediment deposition rates in terms of impact zones (**Figure B7.5.4.2c**), the thresholds for each zone have been assigned as follows (refer **Chapter B3, Coastal Processes** and **Appendix D4, Water Quality Model Development and Calibration Report** for justification):

- **Zone of High Impact** – 1500-mg/cm<sup>2</sup> or more from the cumulative deposition for the capital dredging base case (i.e. cumulative total of 15 mm or greater over the duration of the base case dredge campaign)
- **Zone of Low to Moderate Impact** – Between 500 mg/cm<sup>2</sup> and 1500 mg/cm<sup>2</sup> from the cumulative deposition for the capital dredging base case (i.e. cumulative total of five to 15 mm over the duration of the base case dredge campaign)
- **Zone of Influence (not displayed)** – Less than the 500 mg/cm<sup>2</sup> contour from the cumulative deposition for the capital dredging base case (cumulative total less than five mm over the duration of the base case dredge campaign).

These values are considered conservative, given similar previous approaches elsewhere have used higher thresholds for receptors considered to be highly sensitive to deposition. For example, DHI (2010) use a threshold of >17.5 mm/14 day for likely impacts to corals, and 3.5 to 17.5 mm/14 day for possible impacts to corals. Note that for cumulative sediment deposition, the depths reported for this impact assessment will accrue incrementally over the 34-week duration of the dredging campaign.

As discussed in **Chapter B3, Coastal Processes**, much of the study area is largely a depositional environment. This is particularly true for Trinity Inlet which lacks major fluvial (riverine) flows to drive scour, and is somewhat protected from wave activity. Rather, near shore sediment transport processes throughout the broader study area are largely driven by inputs from the Barron River and associated interactions with wave action along the more exposed coastline to the north of Cairns harbour. Further offshore in the vicinity of the DMPA, marine beds are stable and exhibit little re-suspension. From an ecological perspective, this indicates that marine habitats are prone to sediment deposition near shore, while offshore sedimentary habitats are generally more stable.

### Benthic Fauna

Most infauna fauna species are capable of burrowing at least short distances through sediments, and therefore have low sensitivity to low levels of sediment deposition. Some highly localised smothering of fauna with limited locomotory capacity could occur directly adjacent to the dredge site and DMPA. However for most species, the rates of deposition would be minor compared to the ability of fauna to move through sediments, particularly when deposition is considered incrementally over the duration of the dredging program.

Low to medium density epifauna communities have been recorded on soft sediment habitats within the predicted moderate to high impact zones (**Section B7.4.6**). These communities are comprised mainly of filter feeding fauna (e.g. sea pens, feather stars, sponges) and fauna that entrap their prey (some soft corals). At sub-lethal levels of suspended sediment concentrations, some filter-feeders may benefit from the larger amount of suspended organic matter (i.e. food resources) contained within the dredged material, or released from benthic substrates disturbed by the dredger. It is unlikely that suspended sediment concentrations will reach levels that lead to interference or blocking of the respiratory and feeding structures of these animals. For minor, sub-lethal rates of deposition, many filter feeding fauna are also able to actively self-clean parts of their body prone to trapping unwanted non-food particles. If individuals of these sessile epifauna are very small (recently settled juveniles less than the cumulative deposition depth for a given area, say 15 mm in the Zone of High Impact) they could be smothered, which could lead to stress or mortality. As much of the area within the predicted impact zones is considered to be representative of stable to depositional environments, it is expected that most benthic fauna would be well adapted for coping within the sedimentation rates predicted. The exception to this would be any very small bodied sessile fauna occurring within a Zone of High Impact.

With the exception of artificial hard substrata (sea walls) in the vicinity of the port and Cairns Esplanade, no reefal habitats or reef associated communities are located within areas predicted to be exposed to detectable sedimentation (i.e. outside zone of influence). For intertidal areas, sedimentation is expected to be negligible relative to background levels and natural short-term disturbances such as storm events. Thus, major impacts to intertidal benthic fauna communities are not anticipated.

Based on the predicted low to moderate sedimentation rates over the duration of the dredge program, together with the density and composition of existing benthic communities, sedimentation is expected to result in minor impacts to benthic communities. Most of this impact would be confined to locations within the Zone of High Impact and, to a lesser extent the Zone of Moderate Impact. Note that where sediment deposition is expected to occur in close proximity to areas that are already currently dredged (existing outer channel), inhabitant fauna are likely already exposed to comparable sedimentation rates from ambient hydrological, coastal process and periodic existing maintenance dredging works.

## Flora

Seagrasses (while currently sparse) represent the benthic flora most prone to sedimentation effects. For areas where detectable sediment deposition originates from plumes at the DMPA (i.e. offshore deposition), no notable seagrass beds are known to occur (seagrass was not observed during surveys undertaken for this EIS; if present, it is likely only in sparse isolated patches). As described previously (**Section B7.5.3.1**), sparse isolated patches of deep water seagrasses may occur from time to time. If seagrass occurs in these offshore sediment deposition areas, any burial or smothering of these plants would be considered to be a minor impact. This is based on: i) the anticipated sparse and ephemeral characteristics of any such seagrass (i.e. not critical habitat compared to the more extensive seagrass beds known to occur elsewhere in the study area during favourable conditions); and ii) the temporary and short-term nature of any such effects.

In the current Cairns context, more critical to potential seagrass impacts is the consideration of inshore sediment deposition. Historically, the most extensive meadows tend to be located around the intertidal and shallow subtidal areas of Cairns harbour (**Section B7.4.4**). However, seagrass in these areas is currently in poor condition. It is therefore assumed that seagrass at these inshore areas is presently in the initial stages of recovery, and may be more sensitive to deposition than established meadows. For example, any small-sized shoots and new seedlings occurring during periods of increased sediment deposition may be more prone to smothering and burial in comparison to more mature or established plants. Based on the extent of predicted near shore sediment deposition impacts (refer **Figure B7.5.4.2c**), any potential impacts (if seagrass is present during the dredging campaign) would be expected to be restricted to: i) areas within and adjoining the southern end of the outer channel, seaward from Cairns Esplanade; and ii) parts of Trinity Inlet, along the northern side of Admiralty Island. The risk associated with any such impact would largely depend on the likelihood of seagrass occurring within areas affected by detectable deposition during the dredging activities.

Note that for both offshore and near shore seagrasses, it can be difficult to determine the degree to which sedimentation alone contributes to physiological stress. Sedimentation often occurs in areas also (and simultaneously) impacted by elevated turbidity. This means that when combined with elevated turbidity levels, sediment deposition could result in cumulative stress to seagrasses, particularly during periods when seagrasses are less likely to adapt to sedimentation rates (e.g. autumn-winter). Similar to reduced light levels, this stress could result in localised mortality of existing established seagrasses, or a range of other community changes symptomatic of stress (reduced shoot height, above ground biomass, etc.). Any such effects, however, would only apply to locations where seagrass extent is coincident with sedimentation impact zones at the time of dredging (possibly adjacent to the inshore part of the navigation channel, not applicable to DMPA)

As minor sedimentation impacts are recognised within Trinity Inlet, consideration is also given to potential sedimentation effects to mangroves which can be intolerant of high levels of sedimentation. Natural sedimentation rates within mangrove forests vary spatially and temporally. Elsewhere, they were reported by Ellison (1998) to be up to 10 mm/annum. Natural sedimentation rates for mangroves within the study area, particularly Trinity Inlet, are likely to be much higher than this, given it is an estuary with a naturally a high accretion rate. Excess levels of sedimentation can cause stress to mangroves as a result of smothering and burial of root systems. For many mangrove species, roots provide a critical metabolic function, which can be impaired by burial. This can lead to reduced vigour and mortality, depending on the amount and type of sedimentation, and the species under consideration.

Mangroves present within the study area, particularly Trinity Inlet, are assumed to be well adapted to the local sediment accretion conditions. This, together with the low rates and levels of sedimentation predicted near mangrove areas, especially relative to background levels, suggests that any such effects would be expected to have negligible impacts, if they occur at all.

**Figure B7.5.4.2a Predicted Cumulative Sediment Deposition for the Capital Dredging Base Case (Scale: 50 to 1000)**






Figure B7.5.4.2b Predicted Cumulative Sediment Deposition Occurring after the Capital Dredging Base Case and 12 Months Re-suspension (Scale: 50 to 1000)





Figure B7.5.4.2c Zone of Impact Map for Sediment Deposition



### B7.5.4.3 Maintenance Dredging

Future maintenance dredging will be required to ensure that the dredge footprint remains at the required depths for safe navigation of ships. Compared to capital dredging, much smaller volumes of material are involved in maintenance dredging and the timeframes over which dredging will occur will be shorter. At the outer channel an increase in annual maintenance dredging volume in the order of 80,000-100,000 m<sup>3</sup> per year is expected, while the existing annual maintenance dredging volume for the inner port is not likely to change significantly (refer **Chapter B3, Coastal Processes**). Current channel maintenance dredging campaigns typically occur during the months of July to October and generally take about three-four weeks to complete. In some cases following large wet seasons split campaigns are undertaken with an earlier one-two weeks dredging in May-June. The additional volume associated with the expanded channel will likely extend these campaigns to a period of four-five weeks.

The frequency and duration of turbidity impacts from future maintenance are likely to be similar in nature to those presented in this EIS; albeit occurring over a much smaller duration each year, which limits the amount of material available for re-suspension. **Chapter B5, Marine Water Quality** details potential impacts to water quality from maintenance dredging, utilising actual water quality monitoring data collected over a 12-month period for this EIS (including during annual maintenance dredging in 2013). Impacts from maintenance dredging are considered to be localised and relatively short term with limited increases in turbidity adjacent to sensitive environments. Furthermore, impacts on sensitive receptors from maintenance dredging has been assessed previously (Environment North, 2005 and Worley Parsons, 2010) and considered acceptable to regulatory agencies (as outlined in the Ports North 10 year maintenance dredging permit and LTMP).

Based on the assessment presented in **Chapter B5, Marine Water Quality**, turbid plumes from future maintenance dredging are considered to pose a minor impact to marine water quality. Flow on effects to flora, fauna and other marine ecology values, as a result of both water quality and sediment deposition effects, are likewise considered to minor. Relative to existing maintenance dredging operations, this minor classification takes into account i) the slightly longer duration of future maintenance dredging, and ii) the placement of dredged material at a new DMPA location (as opposed to the existing DMPA). Refer to **Sections B7.5.4.1** and **B7.5.4.2** for discussion on marine ecology impacts associated with dredged material placement at the DMPA, noting that any such effects will be for a shorter duration during future maintenance dredging campaigns.

### B7.5.4.4 Mitigation and Residual Effects

The project design and Dredge Management Plan (**Chapter C2, Dredge Management Plan**) incorporate numerous mitigation measures that will be adopted to reduce the extent and magnitude of turbid plumes in order to minimise impacts to marine flora and fauna. Most significantly, the base case dredging scenario incorporates zero overflow from the TSHD; and the alternative case, if used, would drastically reduce the duration of the dredging program. Specific relevant strategies and components of the DMP include:

- Capital dredging will not be carried out in late spring and summer (November to February). This coincides, in part, with when seagrasses and other benthic biota may be most likely to be undertaking important life history functions (e.g. seagrass growth, coral spawning, spawning on many commercially significant fisheries species) or coping with seasonal environmental stress (e.g. flood-related effects)
- An environmental valve (“green valve”) will be used in overflow pipes of the TSHD to reduce the dispersion of sediments from dredging
- Overflow levels will be raised to the highest allowable point during sailing from the dredge area to the DMPA to ensure spillage of sediment is minimised
- Sailing routes will be optimised to minimise the generation of propeller wash (noting that propeller wash from the dredger has been taken into account in the modelling as a contributor to sedimentation impacts)
- A reactive water quality monitoring program will be developed and implemented. Dredging activities will be modified or suspended in the event that monitoring detects exceedance/s of trigger values, which will illicit various management responses. Water quality baseline data collected for this project (**Chapter 5, Marine Water Quality**), together with local photosynthetic active radiation (PAR) data collected by James Cook University, will be used as the basis for establishing these trigger values
- A seagrass monitoring program (and possibly soft sediment benthos monitoring) will be developed and implemented to identify any changes to communities as a result of the dredging program. This will include sampling at multiple times before and dredging at putative impact sites located adjacent to the near shore project footprint and DMPA, and at suitable control sites.

Overall, the effects of turbidity plumes and sediment deposition are considered to be of Minor significance. Given that such effects are likely to occur, albeit at localised scales, this results in residual medium impact.

## B7.5.5 Interactions between Marine Fauna and Vessels or Dredge Plant

This section examines the potential for interactions between marine fauna and vessels or dredge plant, and associated ecological effects. While invertebrates and fish are mentioned, discussion focuses on marine megafauna species, particularly threatened or otherwise listed species of conservation significance (MNES).

For capital dredging, it is anticipated that the following dredge plant will be required:

- At least one medium-sized trailing suction hopper dredge (TSHD)
- Backhoe dredger and bed leveller
- Work boats/survey boat (discussed below in next section).

Interactions between such vessels and marine fauna may arise during capital dredging or maintenance dredging by way of one or more of the following mechanisms:

- Direct contact or obstruction of fauna passage
- Emissions of artificial noise from the dredger
- Entrainment of fauna at the dredge head
- Emissions of artificial light during night dredging works.

The ongoing operation of the wharf will facilitate an increase in the size and frequency of ship traffic to the CCLT, which will increase the underwater noise and artificial lighting sourced from such vessels, and the potential for direct interactions with marine fauna. Note that such vessels frequent the broader study area but that larger cruise ships are unable to approach the existing port facilities. Rather, large cruise ships currently moor off Yorkeys Knob and ferry passengers by tender to the mainland. This means that impacts relating to interactions between these vessels and marine fauna, essentially relate to a shift in the location of such interactions (i.e. from Yorkeys Knob to the outer channel and inner port), as well as any anticipated increase in cruise ship movements.

### B7.5.5.1 Direct Interactions between Marine Fauna and Vessels or Dredge Plant

When operating any kind of vessel in marine waters, there is a potential risk of fauna vessel strike, primarily for mobile megafauna that swim near the surface and/or frequent the surface to breathe, such as whales, dolphins, dugongs and turtles. Interactions may also occur if the presence of a vessel obstructs fauna passage, which may occur if the presence of a vessel deters an animal from continuing along an intended path of passage, or is inclined to detour significantly around a vessel to reach an intended destination (i.e. avoidance behaviour – discussed further below with respect to potential noise effects).

Large vessels currently operate within the study area, including, the TSHD dredger for annual maintenance dredging and large cruise vessels which moor approximately four km offshore from Yorkeys Knob. Smaller vessel movements (commercial charters, recreational, small cruise vessels) regularly occur throughout the study area. The large vessels specifically associated with this project would represent a small proportion of the total number of boat movements expected to occur over the duration of capital or operational works. However, while large vessels are slow-moving and provide marine fauna time to evade the approaching vessel, they also typically have large powerful propellers and a lower draft. This means that if an animal does not move out of the vessel path, there is a greater risk of severe injury or mortality. In the event that such interactions occur, they would generally occur at locations within the footprint and associated vessel movement paths (between the outer channel and DMPA, offshore shipping route).

Given marine megafauna prone to vessel strike (turtles, dugongs, cetaceans) occur within the study area (refer **Section B7.4.7**), and that these fauna are afforded a high conservation value (MNES), the unmitigated significance of such impacts would be considered to be high. Relevant mitigation measures and the residual risk are described in **Section B7.5.5.3**.

In terms of entrainment, it is possible for the suction at the dredge head to entrain fauna, potentially resulting in fauna injury or mortality. Of the marine megafauna, turtles are the group most likely to be affected by this process. Turtles are highly mobile and will tend to avoid the dredger, typically returning to the surface to breathe every few minutes. However, they can remain underwater for as long as two hours without breathing when they are resting. Dr Col Limpus suggests that sea turtles can use shipping channels as resting or shelter areas, and that there are recorded incidences of turtles being injured by trailer suction hopper type dredgers. GHD (2005), citing personal communication from Dr Limpus, suggest that the numbers of turtles captured during dredging across all Queensland Ports is decreasing, with an average of 1.7 loggerhead turtles per year. Furthermore, it was suggested that current research indicates the impact of dredging on the overall viability of turtle populations is very low compared to the numbers killed by vessel strikes, trawling, fishing, ingestion of marine debris and indigenous hunting. No incidents between dredgers and turtles have been reported in Cairns for the past 10 years, and the potential for such is considered very low, yet possible over the full life of the project. Maintenance dredging with the TSHD *Brisbane* over this period has incorporated turtle deflection devices and the exclusion practices noted below.

Given the relatively low numbers of turtles impacted by dredgers compared to other activities, and the use of effective management and operational practices to reduce the potential for turtle capture, it is not considered that the proposed dredging will have a significant impact on turtle populations in the study area. Other megafauna species (e.g. cetaceans, dugong, etc.) are not considered to be prone to dredge entrainment and will not be impacted by such interactions. Best practice dredging equipment, techniques and management will be used to further reduce risks to turtles (see below).

### **B7.5.5.2 Indirect Interactions between Marine Fauna and Vessels**

#### Lighting

When vessels are operated at night, their on-board lighting systems will generate light emissions to the marine environment. Marine turtles are particularly sensitive to artificial lighting as they may become disorientated during nesting and hatching (Witherington 1992). However, no turtle nesting areas exist in close proximity to the dredge operations or at the inner port, and there is a low incidence of turtle nesting elsewhere in the harbour (i.e. in sight of the outer channel). Further, in the unlikely event that light from project-related vessels can be detected by emerging hatchlings during either project construction or operation, the seaward position of such vessels at all times does not pose a risk for guiding hatchlings landward. In this respect, land-derived lighting such as those at the port or throughout Cairns City presents a significantly greater lighting attraction risk for marine fauna.

Artificial light is not known to have a major effect on foraging patterns of turtles, dolphins or dugongs. Given marine fauna with the potential to be detrimentally affected by lighting (i.e. sea turtles) occur in the study area, and that these fauna are afforded a high conservation value, the significance of artificial lighting affecting these fauna is considered to be a potential Minor Impact. Mitigation strategies will be implemented to further reduce potential impacts (see below).

#### Noise

The production and reception of particular sounds are important to many marine fauna species, particularly marine mammals. Both natural and anthropogenic sounds have the potential to interfere with various biological functions. During construction, noise generated by dredging has the potential to adversely affect megafauna as it will form a persistent source of underwater noise that will continue (intermittently) for the duration of dredging works. Such noise may be generated by mechanical means (vessels engines, dredge gear, propellers and other machinery), or by water movements on the vessel hull. While dredger generated noise is normally unlikely to occur at levels that could cause acute hearing damage to marine fauna, it may cause subtle but possibly more widespread increases to ambient noise levels. This may include for example, masking of biologically important sounds (e.g. vocalisations), interfere with dolphin sonar signals or alter fauna behaviour (i.e. noise avoidance). Similar such effects can be expected during the operational phase of the project when large cruise vessels are approaching or docking at port, or during maintenance dredging campaigns.

In general, the most likely impact of underwater noise from project-associated vessels for marine megafauna is the temporary avoidance of the offending vessels and their immediate surrounds. The inner port is known to support inshore dolphin species, while the wider harbour area supports sea turtles, dolphins, whales, dugongs and other threatened and/or migratory marine species at various times; depending on temporal factors such as migration seasons, availability of food resources, etc. The likelihood of acoustic impacts to marine fauna occurring would depend on whether these fauna are present at the time of vessel operation, the number of animals present and their proximity to the underwater noise source.

If present in or near the dredge footprint during dredge or cruise vessel operation, turtles may exhibit a different response to noise than marine mammals. Turtles often remain stationary for long periods, feeding and resting. GHD (2011) observed turtles exhibiting negligible response in close proximity to marine piling operations and, based on this observation, suggested that it cannot be assumed that turtles will voluntarily move away from adverse vessel/dredge noise effects.

**Chapter B10, Noise and Vibration** details the ambient underwater noise conditions of the study area and associated impacts predicted to result from construction and operations during the life of the project. While information on the effects of noise on marine fauna in an Australian context is extremely limited, dredging is predicted to have negligible impact on marine mammals and dugongs, primarily restricted to localised behavioural changes within ~100-200 m of the dredge (**Chapter B10, Noise and Vibration**). Hearing damage would only be expected if animals remain in the vicinity (~10 m of the dredge) for prolonged periods, which is considered extremely unlikely to occur. For underwater noise associated with shipping, effects to marine fauna are expected to be negligible since similar noise sources currently occur in the study area and localised behavioural changes (avoidance) are considered the most likely effect (refer **Chapter B10, Noise and Vibration**).

As discussed below, mitigation measures will be implemented to further reduce the risk of vessel related noise effects.

### B7.5.5.3 Mitigation and Residual Effects

Management strategies primarily focus on dredge vessels and dredge plant, and will be implemented throughout the course of the proposed capital and operational dredging works to minimise the risk of interactions with vessels. These management strategies are set out in the Dredge Management Plan (**Chapter C2, Dredge Management Plan**) and will include:

- Implementation of marine megafauna management strategies
- Implementation of megafauna exclusion zones (maintaining a given buffer distance between the dredge vessels and megafauna) and associated reactive megafauna monitoring program (regular visual inspections of dredge footprint area and dredge path)
- If visual monitoring for megafauna from the dredge vessels detects megafauna within or headed towards exclusion zones, execute strategies to avoid interactions as required – stopping work if megafauna, especially whales, are within or near exclusion zones; halt dredge vessel transit if there is a likelihood the vessel would encroach on observed whales or their anticipated path
- Operational procedures to minimise the risk of capture of turtles lying on the seabed, especially utilising tickler chains on the dredge head as a fauna exclusion device to reduce fauna entrainment and prevent fauna injury and mortality
- Ensure dredge suction is not started until the dredge head is lowered and in contact with the seafloor, and stops before lifting the dredge head from the seabed
- Where it does not conflict with security and safety requirements, lighting on the dredge vessel will aim for low wattage and/or directional light fixtures.

Together, these mitigation strategies will reduce the likelihood of interactions between the dredge vessels and marine megafauna, such that the overall residual impacts to marine megafauna are considered highly unlikely for all related mechanisms (i.e. vessel strike, noise, entrainment and light).

## B7.5.6 Acoustic Effects to Marine Fauna from Wharf Upgrade Works

### B7.5.6.1 Underwater Noise during Wharf Upgrade

For the wharf upgrade works, underwater noise will result from multiple sources, of which the most chronic will be repeated pulsed inputs from driving the 84 racking steel piles during construction. Noise will also be generated directly from general wharf construction work, as well as through the operation of construction vessels. It is envisaged that the piles will be driven from a barge, by a piling rig with crane and hammer.

**Section B7.5.5.2** (above) describes the general impacts that could occur from interactions between marine fauna and anthropogenic underwater noise. Fish deaths have also occasionally been reported in close proximity to piling as physical damage can occur to non-auditory tissue (e.g. vascular tissue) or air-filled cavities such as swim bladders. As described **Chapter B10, Noise and Vibration**, piling activities during the wharf upgrade works are predicted to have minor impacts on fish, with localised fish mortality within the immediate vicinity (~one-three m) of the piling rig and behavioural changes (avoidance) expected at distances within one km of the piling rig.

Piling noise is expected to have negligible impacts on marine mammals, with the potential for hearing damage limited to the immediate vicinity of the piling rig (up to ~10 m). Although behavioural changes (avoidance) are expected, these are predicted to be limited to within up to ~500 m of the piling rig (**Chapter B10, Noise and Vibration**). Overall, any piling related noise effects to marine megafauna are considered to be temporary (for the duration of construction works) and minimal in the context of the existing noise regime of the area.

### B7.5.6.2 Mitigation and Residual Effects

The following measures, specific to construction phase piling works, will be implemented to reduce underwater noise and associated effects to marine fauna:

- A resilient pad (dolly) will be used where feasible between the pile and hammer head
- A megafauna observation zone of one km, and exclusion zone of 100 m will be adopted, with piling operation shut down if marine megafauna species are observed within or approaching the exclusion zone
- A ‘soft-start’ regime will be adopted at each day’s commencement of piling works
- Underwater noise monitoring will be conducted at the onset of piling to confirm/calibrate the noise predictions, and noise management adapted appropriately.

Implementation of these measures, together with Best Management Practice and Best Available Technology Economically Achievable, is expected to result in negligible, yet possible, impacts to all marine species considered during wharf upgrade works.

## B7.5.7 Other Indirect Interactions between Vessels and Marine Receptors

### B7.5.7.1 Potential for Marine Pest Introductions

There is a risk that dredging plant (construction stage) and other vessels (operational stage) could translocate introduced marine pests from the port of origin to the Port of Cairns. **Section B7.4.8** outlines the few marine pest outbreaks (and eradication operations) that have occurred at the Port of Cairns in recent years. While marine pests, if present, could be transported from the dredge or cruise vessels to the local marine environment, the project is not considered to pose a notable risk in terms of the potential of introducing marine pests to the Cairns harbour and surrounds if appropriate biosecurity inspections and management are employed. This is based on the following:

- The study area is not currently known to support populations of marine pests of concern that could be dispersed by the vessels to waters elsewhere
- The dredge vessel remaining in the study area for the duration of the dredging campaign
- As part of the Dredge Management Plan, appropriate measures will be in place during construction to reduce the potential for introducing marine pests from the dredger (e.g. compliance with antifouling, hull cleaning and ballast treatment requirements)
- The study area is already regularly visited by dredge plant and international cruise ships, in addition to other international vessel traffic
- Cairns is a tropical port and tropical ports are generally less prone to marine pest invasion in comparison to ports in the southern temperate waters of Australia (Hilliard and Raaymakers 1997; Hilliard 1999; Hutchings et al. 2002). This is because i) tropical species often have a widespread equatorial distribution, resulting in a higher probability that Cairns is within their natural geographic range; and ii) foreign temperate species used to cooler waters are less likely to survive and establish in warmer tropical waters
- The study area is not currently known to support populations of marine pests of concern that could be dispersed by the vessels to waters elsewhere.

Notwithstanding this, translocation of exotic marine pests into a new environment is potentially an important issue for the Port of Cairns. The environmental and economic impacts due to the introduction of exotic marine pests can be significant. Marine pests, once established, can be difficult to eradicate and can have serious and permanent consequences for the marine environment, marine productivity and public health. While unlikely, the introduction and subsequent establishment of marine pests would be present a moderate impact from a marine ecology perspective. Accordingly, marine pest risks will be managed in accordance with standard mitigation procedures discussed below.

### B7.5.7.2 Disturbance from Vessel Wash

Wash generated by cruise ships, tugs, dredgers and construction vessels could introduce increased wave energy to intertidal habitats, or lead to the disturbance of the seafloor through propeller wash. In intertidal areas in close proximity to vessel routes (Trinity Inlet, East Trinity) the wash created by vessels could scour intertidal banks and littoral soft sediments, potentially causing adverse alterations to associated flora and fauna communities (mangroves, benthic invertebrates). While these areas currently experience a high level of vessel traffic, the project is expected to result in an increase in the number of large ship movements in the inner port and outer channel (estimated increase in cruise shipping activity of 39 percent by 2016) and, presumably, a corresponding increase in tug activity). This has the potential to increase the magnitude and frequency of adverse vessel wash. A number of mitigation measures may be implemented to reduce these effects (see **Section B7.5.7.3** below) if such impacts become apparent.

Propeller wash could lead to mobilisation of seabed sediments, resulting in scouring in affected areas and possibly disturbance of benthic organisms. It would be expected that the scale of impact of propeller wash to the seabed and associated communities will be negligible, relative to natural hydrological process that prevail in the study area. Note that numerical modelling results presented in **Section B7.5.4** incorporate propeller wash as a source of suspended sediments during dredging.

### B7.5.7.3 Mitigation and Residual Effects

#### Marine Pests

International and domestic vessels involved in either the construction or operational phases of the project will be required to comply with national and state biofouling and ballast water management guidelines, and other requirements to minimise the risk of introductions of marine pest species. Nonetheless, moderate marine pest effects may occur, albeit at an unlikely likelihood.

#### Vessel Wash

Best management practices will be employed to minimise the potential effects of vessel wash to marine habitats. These are outlined in the environmental management plans developed for this project (**Chapter C1, Environmental Management Plan (Construction and Operation)**), of which key relevant elements include:

- Ensuring that construction activities are restricted to the approved development footprint
- Ensuring that vessels operate in designated shipping channels and at designated low speeds to minimise the risk of habitat disturbance from vessel wash.

The residual impact following the implementation of the above measures is classified as Minor.

### B7.5.8 Exposure of Marine Flora and Fauna to Debris, Spills and Dredging-Borne Contaminants

#### B7.5.8.1 Marine Debris

Injury and fatality to vertebrate marine life caused by ingestion of, or entanglement in, harmful marine debris is listed as a key threatening process under the EPBC Act. Construction and operational works will generate large quantities of rubbish which could pose a risk to the marine fauna of the study area if not appropriately managed. In particular, plastic bags and packaging could pose a risk to local marine turtles, whales and fish populations.

A variety of waste management strategies will be employed as part of the environmental management plans developed for this project, to reduce waste generation and the quantity of plastic wastes entering the marine environment (see **Chapter C1, Environmental Management Plan (Construction and Operation)**).

#### B7.5.8.2 Potential Spills

It is possible that chemical spills would occur on, or from the dredger or other vessels such as cruise vessels, creating the potential for dredge-derived potential contaminants to be introduced to the marine environment. These could include, for example, hydrocarbons or other potential toxicants stored on board. Spills could occur either in the vicinity of the dredge footprint, or while vessels are in port or underway outside the project footprint. In the event that a spill occurs, it may present a toxicity risk to marine flora and fauna. The significance of such an impact is highly variable, depending on factors such as:

- The type of material spilt and its chemical constituents
- The volume and/or load concentration of potential toxicants of concern entering the marine environment
- Climate and tidal conditions at the time of event
- The location and timing of a spill, which can dictate the mixing potential (i.e. concentration reduction), extent of water quality effects, and the likelihood of sensitive receptors occurring in the affected area.

Spills of this nature are considered to be unlikely, and no more likely than typical for other large vessels using the wider study area at any given point in time under existing conditions. Given their localised extent or potentially undetectable effects in the event that they do occur, are considered to represent a low level of impact. Implementation of the mitigation measures listed below will further reduce the likelihood of such occurrences to highly unlikely.

#### B7.5.8.3 Potential Toxicants Mobilised by Dredging and Construction

As discussed in **Chapter B4, Marine Sediment Quality**, marine sediments in the proposed dredging and pile driving areas contain potential contaminants at concentrations (95 percent UCL) that are below NAGD screening levels, and therefore do not pose a toxicity threat. As such, impacts from mobilisation of contaminants from the dredging process are expected to be negligible.



Potential acid sulfate soil (PASS) in the marine sediments will be dredged. If exposed to air for extended periods, some acidity would be released from this sediment (refer **Chapter B4, Marine Sediment Quality**). As ocean disposal of the dredged material is proposed, oxidation of the dredge material is considered unlikely since the sediments will stay saturated with seawater. Therefore, the potential impact to water quality, and flow-on effects to marine ecology values, PASS disturbance is considered to be negligible.

Future maintenance dredging and dredge material placement during the operational phase of the project will be undertaken in accordance with NAGD or future versions of these guidelines. As such, the potential impacts associated with this activity are considered to be negligible.

#### **B7.5.8.4 Mitigation and Residual Effects**

Strategies that will be utilised to reduce water quality effects are listed below for:

##### Marine Debris

Throughout both the construction and operational phases of the project, ships, dredgers and other vessels associated with the project will need to ensure waste materials are properly managed in accordance with standard protocols and waste management strategies in the respective Management Plans. The Port of Cairns will provide appropriate waste reception facilities for accommodating this, in line with best practice (e.g. *Best Practice Guidelines for waste Reception Facilities at Ports, Marinas and Boat Harbours in Australia and New Zealand, IMO Guide to Best Practice for Port Reception Facility Providers and Users*).

##### Potential Toxicants Mobilised by Dredging or Construction:

It is assumed that the following standard mitigation measures will be employed to ensure associated impacts are negligible:

- Dredging will be undertaken in a manner consistent with the requirements of the NAGD
- A Dredge Management Plan has been developed for the project (see **Chapter C2, Dredge Management Plan**), which will be implemented throughout the duration of the works. A key component of this plan is a water quality monitoring program that will enable reactive and adaptive management of dredging operations in order to minimise water quality effects and, thus, effects to marine flora and fauna
- Dredge material should remain waterlogged and not be left within TSHD hopper or dump barges for periods longer than 24 hours to minimise the risk of PASS oxidation
- Future maintenance dredging during the operational phase of the project is undertaken consistent with the LTSDMP and in accordance with NAGD or future versions of these guidelines. Also, maintenance dredge material is tested in accordance with the frequency intervals outlined in the guidelines to ensure only acceptable, uncontaminated maintenance dredge material will be placed at sea.

##### Spills

The following additional mitigation is proposed to reduce the potential impacts associated with spills during project construction and operation:

- Hazardous material handling procedures have been developed for the project as part of the Dredge Management Plan (see **Chapter C2, Dredge Management Plan**)
- Emergency spill response procedures will be implemented if/when required
- Relevant staff will be trained to ensure they have an appropriate level of competency for executing the above spills procedures
- Revise fuel handling and spill response procedures in the port's operational procedures to minimise the potential future risk to sediment quality from refuelling activities associated with the provision of IFO at the port.

With the implementation of the above measures, it is considered highly unlikely that spills, if they occur, will cause adverse impacts to the marine environment. Note that spill and emergency response procedures will be outlined as part of the Dredge Management Plan.

Overall, through the implementation of the above strategies, the residual impact to marine ecological values is considered to be Minor and highly unlikely.

## B7.5.9 Summary of Impacts to Marine Megafauna

### B7.5.9.1 Key Impacting Processes Considered for Marine Megafauna

As described in the above sections, the main processes with the potential to impact marine megafauna communities within the study area are:

- Modifications to benthic habitats and associated benthic communities as a i) direct result of dredging in the dredge footprint (**Section B7.5.3**), or ii) through water quality and sedimentation effects resulting from turbid plumes (**Section B7.5.4**)
- Direct interactions between megafauna and vessels or dredge plant (i.e. entrainment or vessel strike) (**Section B7.5.5**)
- Indirect interactions between megafauna and vessels or pile driving works (i.e. artificial lighting and noise) (**Sections B7.5.5** and **B7.5.6**)
- Potential exposure of megafauna to marine debris (**Section B7.5.8**).

In general, the following conclusions are drawn with respect to marine megafauna impacts:

- Turbid plumes generated by either capital or maintenance dredging and dredged material placement are not expected to result in direct effects to marine megafauna. Some species such as green and hawksbill turtle, which feed on seagrass and/or in reef environments, may avoid areas affected by turbid plumes which will vary in the area affected depending upon dredge operational strategy and climatic conditions. However, it is noted that key foraging habitats for these species (i.e. reefs, notable seagrass beds) do not coincide with the predicted extent of turbidity impact zones
- Effects resulting from direct loss or disturbance of benthic habitats are considered unlikely, since benthic habitats within the dredge footprint, DMPA or pile driving footprint are unlikely to provide critical foraging or other habitat functions for green turtles and dugongs (particularly since seagrass in these areas is absent or sparse, relative to elsewhere in the study area). Impacts to other megafauna are likewise considered unlikely as soft sediment habitats, similar to those in the immediate footprint, are widespread throughout the remainder of the study area
- Vessel fauna strike, or entrainment (at the dredge head) may occur, but this is considered a very low probability and is dependent on highly variable factors such as the species present, the number of individuals present and their condition, at the time of such works/operations. Numerous mitigation measures will be implemented to reduce this risk
- Lighting effects to marine megafauna are considered to be negligible, especially near shore in the context of land derived light sources
- Potential noise effects may be caused by either pulsed pile driving works, or the more persistent (albeit temporary) dredge and vessel sources. For both of these, the most likely effect is expected to be temporary megafauna behavioural changes, by way of avoidance of the noise source and its immediate surrounds. Hearing damage would only be expected if animals remain in close range to the noise source for prolonged periods. This is considered unlikely for most marine megafauna.

### B7.5.9.2 Mitigation and Residual Effects

Key mitigation strategies specific to minimising potential harm to marine megafauna as a result of the project are primarily focused on reducing interactions (direct or indirect) between megafauna and vessels, dredge plant and pile driving. These include management strategies set out in the Dredge Management Plan (**Chapter C2, Dredge Management Plan**), as listed in **Section B7.5.5.3** and management strategies to be implemented during pile driving works as listed in **Section B7.5.6.2**. These measures were developed in line with current best industry practice, and also consider general management strategies outlined in species recovery plans (including the *2003 Recovery Plan for Marine Turtles in Australia*).

Mitigation measures focusing on protecting megafauna habitats (e.g. seagrass) are provided elsewhere, in the previous sections of this impact assessment.

Overall, implementation of these mitigation strategies, together with best practice construction and waste management methodologies, are expected to result in minor residual impacts for marine megafauna.

## B7.6 Assessment Summary

A summary of the outcomes of the risk-based assessment for activities that have the potential to impact on the marine ecology values of the study area, during either the construction and/or operational phases of the project, are summarised in **Table B7.6a**. Most potential impact processes are rated as having a negligible to low risk to marine ecology. The exceptions to this are:

- Construction phase habitat modifications, which rated as a medium impact since they will be permanent within the project footprint
- Construction phase dredge plumes, which rated as medium impact, primarily because there is a risk of moderate impacts to seagrass in recovering seagrass areas if turbid plumes coincide with the location of new growth (new shoots, seedlings) and have a negative ecological effect
- The potential for marine pest introductions, rated as a medium impact during the operational phase, which is when a significant increase in international ship traffic is anticipated.

Table B7.6a Assessment Summary Table – Marine Ecology

Marine Ecology Primary impacting processes	Initial assessment with standard mitigation in place			Residual Assessment with additional mitigation in place				
	Statutory mitigation measures required	Significance of impact	Likelihood of impact	Risk rating	Additional mitigation measures proposed	Significance of impact	Likelihood of impact	Risk rating
<b>CONSTRUCTION PHASE</b>								
Direct modification of benthic habitats (subtidal soft sediment habitats) from capital dredging, dredged material placement and wharf upgrade works	<ul style="list-style-type: none"> <li>Project design minimises area of marine habitat directly lost/affected by proposal.</li> <li>Selection of preferred DMIPA option takes minimising marine ecological effects into account.</li> </ul>	Moderate	Likely	Medium	Implementation of DMP, including <ul style="list-style-type: none"> <li>Reactive management following operational bathymetric surveys.</li> </ul>	Minor	Likely	Medium
Increased suspended sediment concentrations from capital dredging and dredged material placement (resultant water quality and sediment deposition effects)	<ul style="list-style-type: none"> <li>Project design minimises the extent (volume) of dredging</li> <li>Ensure TSHD dredge operates within the approved dredge footprint at all times</li> <li>Overflow dredging by the TSHD is not undertaken unless as required by and in accordance with the worst case (limited dredging) scenario.</li> <li>Dredge hopper compartment is to be kept water tight during all dredging activities, except placement at the DMIPA</li> <li>Ensure the top of overflow valves are not lowered during the transport component of the dredging cycle</li> <li>No high pressure jets to be used on drag heads outside of DMIPA</li> <li>Dredge to be fitted with a 'green valve'.</li> </ul>	<p><u>Seagrass:</u> recovering / new growth - Moderate</p> <p><u>Seagrass:</u> existing / established meadow - Minor</p> <p><u>Other receptors:</u> coral reefs, megafauna, fisheries - Negligible</p>	<p>Seagrass: recovering / new growth - Likely</p> <p>Seagrass: existing / established meadow - Possible</p> <p>Other receptors: coral reefs, megafauna, fisheries - Possible</p>	Medium	Implementation of DMP, including <ul style="list-style-type: none"> <li>avoidance of summer months for dredging</li> <li>sailing routes optimised to minimise propeller wash</li> <li>implementation of reactive water quality monitoring program</li> <li>implementation of soft sediment benthos and seagrass monitoring program.</li> </ul>	<p><u>Seagrass:</u> recovering / new growth - Moderate</p> <p><u>Seagrass:</u> existing / established meadow - Minor</p> <p><u>Other receptors:</u> coral reefs, megafauna, fisheries - Negligible</p>	<p><u>Seagrass:</u> recovering / new growth - Possible</p> <p><u>Seagrass:</u> existing / established meadow - Unlikely</p> <p><u>Other receptors:</u> coral reefs, megafauna, fisheries - Unlikely</p>	Medium

<p>Interactions between marine fauna and vessels or dredge plant (i.e. fauna vessel strike, entrainment, indirect artificial noise and lighting effects)</p>	<p>None identified</p>	<p>Moderate</p>	<p>Possible</p>	<p>Medium</p>	<p>Implementation of DMP, including:</p> <ul style="list-style-type: none"> <li>• implementation of marine megafauna management strategies</li> <li>• implementation of megafauna exclusions zones</li> <li>• visual monitoring and implementation of reactive strategies</li> <li>• utilisation of tickler chains and similar dredge device(s) to reduced entrainment risk</li> <li>• cease dredge suction prior to lifting dredge head from seabed</li> <li>• minimise lighting utilisation as practicable.</li> </ul>	<p>Moderate</p>	<p>Highly unlikely</p>	<p>Low</p>
<p>Acoustic effects to marine fauna from wharf upgrade works (especially pile driving)</p>	<ul style="list-style-type: none"> <li>• Implementation of Best Management Practice</li> <li>• Best Available Technology Economically Achievable.</li> </ul>	<p>Minor</p>	<p>Likely</p>	<p>Medium</p>	<ul style="list-style-type: none"> <li>• Resilient pad (dolly) used where feasible between the pile and hammer head</li> <li>• A megafauna observation zone of one km, and exclusion zone of 100 m will be adopted, with piling operation shut down if marine megafauna species are observed within or approaching the exclusion zone</li> <li>• A 'soft-start'/'ramp-up' regime will be adopted at each day's commencement of piling works</li> <li>• Underwater noise monitoring conducted at the onset of piling to confirm/calibrate the noise predictions, and noise management adapted appropriately.</li> </ul>	<p>Negligible</p>	<p>Possible</p>	<p>Negligible</p>

Other indirect interactions from vessels: marine pests introductions	<ul style="list-style-type: none"> <li>• Vessel compliance with national and state biofouling and ballast water management procedures.</li> </ul>	Moderate	Unlikely	Low	None identified	Moderate	Unlikely	Low
Other indirect interactions from vessels: vessel wash	None identified	Minor	Possible	Low	Implementation of construction EMP, including: <ul style="list-style-type: none"> <li>• Ensuring construction activities are restricted to the project footprint</li> <li>• Ensure vessels operate in designated shipping channels</li> <li>• Implementation of go slow zones to minimise wash</li> </ul>	Minor	Unlikely	Low
Introduction of marine debris	<ul style="list-style-type: none"> <li>• Port of Cairns to provide appropriate waste reception facilities, in line with best practice</li> <li>• All vessels to comply with standard waste management protocols.</li> </ul>	Minor	Unlikely	Low	<ul style="list-style-type: none"> <li>• Implementation of waste management strategies in construction EMP to reduce waste generation and the quantity of plastics entering the marine environment.</li> </ul>	Minor	Highly unlikely	Negligible
Risk of toxic spills from vessels	None identified	Minor	Possible	Low	<ul style="list-style-type: none"> <li>• Implement hazardous material handling procedures</li> <li>• Implement emergency spill response procedures as required</li> <li>• Ensure relevant staff trained in above procedures to ensure competency</li> <li>• Revise port fuel handling and emergency response procedures.</li> </ul>	Minor	Highly unlikely	Negligible

Potential contaminants mobilised by dredging	<ul style="list-style-type: none"> <li>• Completion of sediment sampling and analysis program (SAP) in line with NAGD guidelines to characterise sediments and determine suitability for ocean disposal</li> <li>• Based on SAP outcomes, ensure only acceptable material destined for DMPA.</li> </ul>	Negligible	Possible	Negligible	Implementation of DMP, including: <ul style="list-style-type: none"> <li>• Reactive water quality monitoring program and relevant adaptive management strategies</li> <li>• Dredged material to remain waterlogged and not to remain in TSHD hopper or dump barge for periods exceeding 24 hours.</li> </ul>	Negligible	Highly unlikely	Negligible	
<b>OPERATIONAL PHASE</b>									
Increased suspended sediment concentrations from maintenance dredging and dredged material placement (resultant water quality and sediment deposition effects)	<ul style="list-style-type: none"> <li>• Ensure TSHD dredge operates within the approved dredge footprint at all times</li> <li>• Dredge hopper compartment is to be kept water tight during all dredging activities, except placement at the DMPA</li> <li>• Ensure the top of overflow valves are not lowered during the transport component of the dredging cycle</li> <li>• No high pressure jets to be used on drag heads outside of DMPA</li> <li>• Dredge to be fitted with a 'green valve'</li> </ul>	Minor	Likely	Medium	Implementation of DMP, including: <ul style="list-style-type: none"> <li>• sailing routes optimised to minimise propeller wash</li> <li>• implementation of reactive water quality monitoring program</li> <li>• implementation of soft sediment benthos and seagrass monitoring program.</li> </ul>	Minor	Unlikely	Low	
Interactions between marine fauna and vessels/ships or dredge plant (i.e. fauna vessel strike, entrainment, indirect artificial noise and lighting effects)	None identified	Noderate	Possible	Medium	For maintenance dredging, implementation of DMP, including: <ul style="list-style-type: none"> <li>• implementation of marine megafauna management strategies</li> <li>• implementation of megafauna exclusions zones</li> </ul>	Moderate	Highly unlikely	Low	

Other indirect interactions from vessels: marine pests introductions	<ul style="list-style-type: none"> <li>Vessel compliance with national and state biofouling and ballast water management procedures.</li> </ul>	Moderate	Possible (higher frequency of international ships compared with construction phase)	Medium	None identified	<ul style="list-style-type: none"> <li>visual monitoring and implementation of reactive strategies</li> <li>utilisation of tickler chains and similar dredge device(s) to reduce entrainment risk</li> <li>cease dredge suction prior to lifting dredge head from seabed</li> <li>minimise lighting utilisation as practicable</li> </ul>	Moderate	Possible	Medium
Other indirect interactions from vessels: vessel wash	None identified	Minor	Possible	Low	Implementation of construction EMP, including: <ul style="list-style-type: none"> <li>Ensuring construction activities are restricted to the project footprint</li> <li>Ensure vessels operate in designated shipping channels</li> <li>Implementation of go slow zones to minimise wash.</li> </ul>	Minor	Unlikely	Low	
Introduction of marine debris	<ul style="list-style-type: none"> <li>Port of Cairns to provide appropriate waste reception facilities, in line with best practice</li> <li>All vessels to comply with standard waste management protocols</li> </ul>	Minor	Unlikely	Low	<ul style="list-style-type: none"> <li>Implementation of waste management strategies in operational EMP to reduce waste generation and the quantity of plastics entering the marine environment.</li> </ul>	Minor	Highly Unlikely	Negligible	



Risk of toxic spills from vessels	None identified	Minor	Possible	Low	<ul style="list-style-type: none"> <li>Implement hazardous material handling procedures</li> <li>Implement emergency spill response procedures as required</li> <li>Ensure relevant staff trained in above procedures to ensure competency</li> <li>Revise port fuel handling and emergency response procedures.</li> </ul>	Minor	Highly Unlikely	Negligible
Potential contaminants mobilised by maintenance dredging	<ul style="list-style-type: none"> <li>Completion of sediment sampling and analysis program (SAP) in line with NAGD guidelines to characterise sediments and determine suitability for ocean disposal</li> <li>Based on SAP outcomes, ensure only acceptable material destined for DMPA.</li> </ul>	Negligible	Possible	Negligible	Implementation of DMP, including: <ul style="list-style-type: none"> <li>Reactive water quality monitoring program and relevant adaptive management strategies</li> <li>Dredged material to remain waterlogged and not to remain in TSHD hopper or dump barge for periods exceeding 24 hours.</li> </ul>	Negligible	Highly Unlikely	Negligible

## B7.7 References

- Accad, A. and Neil, D. (2006) Modelling pre-clearing vegetation distribution using GIS-integrated statistical, ecological and data models: A case study from the wet tropics of north eastern Australia. *Ecological Modelling* 198 (1-2): 85-100.
- Anthony, K., & Larcombe, P. (2000). Coral reefs in turbid waters: sediment-induced stresses in corals and likely mechanisms of adaptation. In Proc. 9th Int. Coral Reef Symp, 239-244.
- Australian Government (2013) Marine pests interactive map. Marine Pests – the national system for the prevention and management of marine pest incursions. Australian Government, Canberra. <http://marinepests.gov.au/> accessed 20 August 2013.
- Bach, S., Borum, J., Fortes, M., & Duarte, C. (1998). Species composition and plant performance of mixed seagrass beds along a siltation gradient at Cape Bolinao, The Philippines. *Marine Ecology Progress Series* 174, 247-256.
- Bell, J. D., and Pollard, D. A. (1989). Ecology of fish assemblages and fisheries associated with seagrasses. In A. W. D. Larkum, A. J. McComb and S. A. Shepherd (Eds.), *Biology of Seagrasses: A treatise on the biology of seagrasses with special reference to the Australian region* (pp. 565-609). Amsterdam: Elsevier.
- Blaber, S.J.M. and Blaber, T. G. (1980) Factors affecting the distribution of juvenile estuarine and inshore fish. *Journal of Fish Biology*. 17: 143-162.
- Brand-Gardner, S.J., Lanyon, J.M., Limpus, C.J. (1999) Diet selection by immature green turtles *Chelonia mydas*, in sub-tropical Moreton Bay, south-east Queensland. *Australian Journal of Zoology* 47: 181-191.
- Campbell, S., C. Miller, Steven, A., Stephens, A. (2003). "Photosynthetic responses of two temperate seagrasses across a water quality gradient using chlorophyll fluorescence." *Journal of Experimental Marine Biology and Ecology* 291(1): 57-78.
- Campbell, S.J. and McKenzie, L.J. (2004) Flood related loss and recovery of intertidal seagrass meadows in southern Queensland. *Australia Estuarine, Coastal and Shelf Science* 60: 477-490.
- Carlton, J. M. (1974). "Land-building and Stabilization by Mangroves." *Environmental Conservation* 1: 285-294.
- Carruthers, T.J.B., Dennison, W.C., Longstaff, B.J., Waycott, M., Abal, E.G., McKenzie, L.J., Lee Long, W.J. (2002). Seagrass habitats of northeast Australia: models of key processes and controls. *Bulletin of Marine Science of the Gulf and Caribbean*, 71(3), 1153-1169.
- Chartrand, K., Rasheed, M. A., & Sankey, T. (2008). Deepwater seagrass dynamics in Hay Point – Measuring variability and monitoring impacts of capital dredging. n.s: Ports Corporation of Queensland.
- Chartrand KM, R. P. (2012). Development of a Light-Based Seagrass Management Approach for the Gladstone Western Basin Dredging Program. Cairns: State of Queensland (Department of Agriculture, Forestry and Fisheries).
- Coles, R. G., Lee Long, W.J., Helmke, S.A., Bennett, R.E., Miller, K.J., Derbyshire, K.J. (1992). Seagrass beds and juvenile prawn and fish nursery grounds: Cairns to Bowen. Queensland, Department of Primary Industries Information Series 64.
- Coles, R. G., W. J. Lee Long, Squire, L.C., (1985). Seagrass beds and prawn nursery grounds between Cape York and Cairns, DPI.
- Coles, R., J. Mellors, Bibby, J., Squire, B. (1987). Seagrass beds and juvenile prawn nursery grounds between Bowen and Water Park Point. Brisbane, Queensland Department of Primary Industries.
- Coles, R., McKenzie, L. and Campbell, S. (2003) The seagrasses of Eastern Australia, in Green, E.P. and Frederick, T.S. (eds), *World Atlas of Seagrasses*, UNEP World Conservation Monitoring Centre, Cambridge.
- Collier, C., Lavery, P., Ralph, P., & Masini, R. (2009). Shade-induced response and recovery of the seagrass *Posidonia sinuosa*. *Journal of Experimental Marine Biology and Ecology* 370, 89-103.

Connolly, R., G. Jenkins, Loneregan, N. (1999). Seagrass dynamics and fisheries sustainability. Seagrass in Australia - Strategic Review and Development of an R & D Plan. A. J. Butler and P. Jernaakoff. Collingwood, CSIRO Publishing: 25-64.

Connolly, R. M. and M. A. Guest (2004). Critical estuarine habitats for foodwebs supporting fisheries in Port Curtis, central Queensland, Australia, Cooperative Research Centre for Coastal Zone, Estuary and Waterway Management.

Cruz-Motta, J., & Collins, J. (2004). Impacts of dredged material disposal on a tropical soft-bottom benthic assemblage. Marine Pollution Bulletin 48, 270-280.

Dall, W., Hill, B., Rothlisberg, P., & Staples, D. (1991). Biology of the Penaeidae; Advances in Marine Biology Series, Vol. 27. London: Academic press.

Davis, T. (1987). Biology of wildstock Lates calcarifer in northern Australia, in Copland, J.W. and Grey, D.L. Management of Wild and Cultured Sea Bass/Barramundi (Lates calcarifer). Canberra: ACIAR Proceedings 20.

DEWHA (2007) Marine Species Conservation: Leatherback turtle (*Dermochelys coriacea*) Accessed: March 2011 Available: [www.environment.gov.au/coasts/species/turtles/leatherback.html](http://www.environment.gov.au/coasts/species/turtles/leatherback.html)

Department of Environment and Resource Management (DERM) (2010) Wildlife and ecosystems: Humpback whales *Megaptera novaeangliae* Accessed: March 2011 Available: [www.derm.qld.gov.au/wildlife-ecosystems/wildlife/az\\_of\\_animals/whales.html](http://www.derm.qld.gov.au/wildlife-ecosystems/wildlife/az_of_animals/whales.html)

DNPRSR (2012) Declared Fish Habitat Area Summary – Trinity Inlet. Online fact sheet available through the Queensland Government Department of National Parks, Recreation, Sport and Racing. <http://www.nprsr.qld.gov.au/managing/habitat-areas/index.html>

DSEWPAC (2013) Draft EPBC Act referral guidelines for the outstanding universal value of the Great Barrier Reef World Heritage Area. Australian Government Department of Sustainability, Environment, Water, Population and Communities, Canberra.

DSEWPAC/GBRMPA (2013) Guidelines for an Environmental Impact statement for the Cairns Shipping Development (Trinity Inlet) Project, in Port of Cairns and Great Barrier Reef Marine Park, Queensland – March 2013. Jointly prepared by the Australian Government Department of Sustainability, Environment, Water, Population and Communities and the Great Barrier Reef Park Marine Authority.

DSITIA (2013) Wildlife Online extract for Cairns Harbour and surrounds (20 km radius). Extracted 5 September 2013. Queensland Government Department of Science, Information Technology, Innovation and the Arts.

Duarte, C., Terrados, J., Agawin, N., Fortes, M., Bach, S., & Kenworthy, W. (1997). Response of a mixed Philippine seagrass meadow to experimental burial. Marine Ecology Progress Series. 147, 285-294.

Edgar, G. and H. Kirkman (1989). Recovery and Restoration of Seagrass Habitat of Significance to Commercial Fisheries, Victorian Institute of Marine Sciences.

Ellison, J. (1998). Impacts of sediment burial on mangroves. Marine Pollution Bulletin 37, 420–426.

Environment North (2005) Cairns Harbour Dredging and Long Term Dredge Spoil Disposal Management Plan. Prepared for Cairns Port Authority by Environment North, in association with Hydrobiology and NIWA Australia, Cairns.

Freeman, A., Short, F., Isnain, I., Razak, F., & Coles, R. (2008). Seagrass on the edge: Land use practices threaten coastal seagrass communities in Sabah, Malaysia. Biological Conservation 141, 2993-3005.

GBRMPA (2012) Sawfish – A Vulnerability Assessment for the Great Barrier Reef. Online information sheet, accessed 10 September 2013.

[http://www.gbrmpa.gov.au/\\_\\_data/assets/pdf\\_file/0016/21742/gbrmpa-VA-Sawfish-11-7-12.pdf](http://www.gbrmpa.gov.au/__data/assets/pdf_file/0016/21742/gbrmpa-VA-Sawfish-11-7-12.pdf)

GHD. (2005). Port of Hay Point Apron Area and Departure Path Capital Dredging Draft Environmental Impact Statement. n.s: GHD, for Ports Corporation of Queensland.

GHD (2009) Townsville Marine Precinct Project – Environmental Impact Statement, Main Report. Prepared for the Port of Townsville Limited, 2009.

GHD. (2011). Port of Townsville Limited Report for Port Expansion - Marine Megafauna. Townsville: GHD, for Port of Townsville Limited.

Gillanders, B. M., and Kingsford, M. J. (2002). Impact of changes in flow of freshwater on estuarine and open coastal habitats and the associated organisms. *Oceanography and Marine Biology: An Annual Review*, 40, 233-309.

Goudkamp, K. and Chin, A. (2006) 'Mangroves and Saltmarshes' in Chin, A. (ed) *The State of the Great Barrier Reef Online*, Great Barrier Reef Marine Park Authority, Townsville. Viewed 02/09/2013: [http://www.gbrmpa.gov.au/publications/sort/mangroves\\_saltmarshes](http://www.gbrmpa.gov.au/publications/sort/mangroves_saltmarshes).

Grice, A., Loneragan, N., & Dennison, W. (1996). Light intensity and the interactions between physiology, morphology and stable isotope ratios in five species of seagrass. *Exp. Mar. Biol. Ecol.* 195, 91–110.

Haywood, M. D. E., Vance D. J., Loneragan, N. R. (1995). "Seagrass and algal beds as nursery habitats for Tiger Prawns (*Penaeus semisulcatus* and *P. esculentus*)." *Marine Biology* 122: 213-223.

Hazel, J., Gyuris, E. (2006) Vessel-related mortality of sea turtles in Queensland, Australia. *Wildlife Research*, 33: 149-154.

Heasman, M. P., D. R. Fielder, Shepherd, R.K. (1985). "Mating and spawning in the mudcrab, *Scylla serrata* (Forskål) (Decapoda: Portunidae), in Moreton Bay, Queensland." *Australian Journal of Marine and Freshwater Research* 36: 773-783.

Helmke, S., Bailey, S., Garrett, R., Maccoll, K. (2003) Fisheries resources in Trinity Inlet, north-eastern Australia. Annual Assessment from 1997 to 2000, Project Report. Queensland Government Department of Primary Industries.

Hewitt, C. and Campbell, M. (2010) The relative contribution of vectors to the introduction and translocation of marine invasive species. Prepared for the Department of Agriculture, Fisheries and Forestry (DAFF).

Hilliard, R. (1999) Marine pest invasion risks – warm versus cool-water ports. *EcoPorts Monogr. Ser. No.19*: 65-70.

Hilliard, R., Raaymakers, S. (1997) Ballast water risk assessment for 12 Queensland ports, Stage 5: Executive Summary and Synthesis of Results. *EcoPorts Monogr. Ser. No.14*.

Hobday, D., Officer, R., & Parry, G. (1999). Changes to demersal fish communities in Port Phillip Bay, Australia, over two decades, 1970-91. *Marine and Freshwater Research*. 50, 397-407.

Hutchings, P. A. and P. Saenger (1987). *Ecology of Mangroves*. St Lucia, University of Queensland Press.

Hutchings, P., Hilliard, R., Coles, S. (2002) Species introductions and potential for marine pest invasions into tropical marine communities, with special reference to the Indo-Pacific. *Pacific Science*, 56: 223-233.

Hydrobiology. (2003). *Monitoring of Benthos, Port of Hay Point*. n.s: Hydrobiology Pty Ltd, for Ports Corporation of Queensland.

IUCN (2010) *The IUCN Red List of Threatened Species (Version 2010.4)*. Accessed: March 2011 Available: [www.iucnredlist.org/apps/redlist/search](http://www.iucnredlist.org/apps/redlist/search)

Jarvis, J.C., McKenna, S.A. and Rasheed, M.A. (2013). *Seagrass Habitat of Cairns Harbour and Trinity Inlet: Sediment Seedbank Analysis*. Report No. 13/48, James Cook University Publication, Centre for Tropical Water and Aquatic Ecosystem Research, Cairns.

Jarvis, J.C., Rasheed, M.A., McKenna, S.A. and Sankey, T. (2014). *Seagrass Habitat of Cairns Harbour and Trinity Inlet: Annual and Quarterly Monitoring Report 2013*. JCU Publication, Centre for Tropical Water and Aquatic Ecosystem Research, Cairns.

- Jelbart, J. E. (2004). The influence of seascape spatial features on the fish and macroinvertebrates in seagrass beds. Hawkesbury, University of Western Sydney: 165.
- Jenkins, G., & McKinnon, L. (2006). Channel Deepening Supplementary Environment Effects Statement – Aquaculture and Fisheries. Internal Report No. 77. Queenscliff: Primary Industries Research Victoria.
- Johnson, J.E., Brando, V.E., Devlin, M.J., Kennedy, K., McKenzie, L., Morris, S., Schaffelke, B., Thompson, A., Waterhouse, J. and Waycott, M. (2011) Reef Rescue Marine Monitoring Program: 2009/2010 Synthesis Report. Report prepared by the Reef and Rainforest Research Centre Consortium of monitoring providers for the Great Barrier Reef Marine Park Authority. Reef and Rainforest Research Centre Limited, Cairns.
- Jones, G., & Candy, S. (1981). Effects of dredging on the macrobenthic infauna of Botany Bay. Australian Journal of Marine and Freshwater Research, 32, 379-398.
- Kailola, P. J., Williams, M. J., Stewart, P. C., Reichelt, R. E., McNee, A., and Grieve, C. (1993). Australian Fisheries Resources. Canberra: Bureau of Resource Sciences, Department of Primary Industries and Energy, and the Fisheries Research and Development Corporation.
- Kenworthy, W. J. (2000). The role of sexual reproduction in maintaining populations of *Halophila decipiens*: implications for the biodiversity and conservation of tropical seagrass ecosystems. Pac. Cons. Biol., 5, 260-268.
- Lee Long W.J., Mellors J.E., Coles R.G. (1993) Seagrasses between Cape York and Hervey Bay, Queensland, Australia. Australian Journal of Marine and Freshwater Research 44: 19-31.
- Lee Long, W. J., L. J. McKenzie, Coles, R.G. (1996). Seagrass Communities in the Shoalwater Bay Region, Queensland, Great Barrier Reef Marine Park Authority.
- Limpus, C.J., Couper, P.J. and Read, M.A. (1994). The loggerhead turtle, *Caretta*, in Queensland: population structure in a warm temperate feeding area. Memoirs of the Queensland Museum. 37:195-204.
- Limpus, C. (2007) A biological review of Australian marine turtles. 5. Flatback turtle *Natator depressus* (Garman). Queensland Government Environment Protection Agency.
- Limpus, C. (2008) A biological review of Australian marine turtles. 2. Green turtle *Chelonia mydas* (Linnaeus). Queensland Government Environment Protection Agency.
- Loneragan, N. R., and Bunn, S. E. (1999). River flows and estuarine ecosystems: Implications for coastal fisheries from a review and a case study of the Logan River, southeast Queensland. Australian Journal of Ecology, 24, 431-440.
- Longstaff, B., Loneragan, N., O'Donohue, M., & Dennison, W. C. (1999). Effects of light deprivation on the survival and recovery of the seagrass *Halophila ovalis* (R.Br.) Hook. Journal of Experimental Marine Biology and Ecology 234, 1-27.
- Ludescher, C.M. (1997) Fisheries Resources between Bowen and Tully - An Inventory. Report prepared for The Townsville Zonal Advisory Committee and Queensland Fisheries Management Authority. James Cook University, Townsville.
- Marsh, H. and Saalfeld, W.K. (1990) The distribution and abundance of dugongs in the Great Barrier Reef Marine Park south of Cape Bedford. Aust. Wildl. Res. 17:511-24.
- Marsh, H., Corkeron, P., Lawler, I., Preen, A., Lanyon, J. (1996) The status of the dugong in the southern Great Barrier Reef Marine Park. James Cook University, Great Barrier Reef Marine Park Authority Research Publication No. 41. Townsville.
- Marsh, H., De'ath, G., Gribble, N., Lane, B. (1999) Shark control records hindcast serious decline in dugong numbers off the urban coast of Queensland. Great Barrier Reef Marine Park Authority Research Publication No. 70, prepared by James Cook University, Townsville.
- Russell, D.J. and McDougall, A.J. and Fletcher, A.S. and Ovenden, J.R. and Street, R. (2003) *Biology, Management and Genetic Stock Structure of Mangrove Jack, (Lutjanus argentimaculatus) in Australia*. Project Report. Department of Primary Industries, Queensland.

McKay, R. (1992). FAO Species Catalogue, Vol 14, Sillaginid fishes of the world (Sillaginidae). An annotated and illustrated catalogue of the Sillago, smelt or Indo-Pacific whiting species known to date. n.s: FAO Fish Synopses 125.

McKenna SA, Rasheed MA, Sankey, T. & Tol S.J. (2013). Benthic macro-invertebrates of Cairns Harbour and Trinity Inlet: baseline survey – 2012/13. JCU Publication, Centre for Tropical Water & Aquatic Ecosystem Research Publication 13/17, Cairns, 33 pp.

McKenzie, L. (1994) Seasonal changes in biomass and shoot characteristics of a *Zostera capricorni* Aschers. Dominant meadow in Cairns Harbour, Northern Queensland. Aust. Journal of Marine and Freshwater Research, 45: 1337-52.

McKenzie, L., Unsworth, R., Waycott, M. (2012) Reef Rescue Marine Monitoring Program – Intertidal seagrass annual report, for sampling period 1 September 2009 – 31 May 2010. Fisheries Queensland, Cairns.

Meager, J. J. (2003). The Microhabitat Distribution of Juvenile Banana Prawns, *Penaeus merguensis* de Man in Subtropical Eastern Australia and Processes Affecting their Distribution and Abundance. (PhD), Queensland University of Technology, Brisbane.

Meager, J.J., Limpus, C.L. (2012) Marine wildlife stranding and mortality database annual report 2011. III. Marine turtle. Conservation Technical Data Report, Queensland Government Department of Environment and Heritage Protection.

Meager, J.J., Limpus, C.L. (2012) Marine wildlife stranding and mortality database annual report 2011. I. Dugong. Conservation Technical Data Report, Queensland Government Department of Environment and Heritage Protection.

Meager, J.J., Winter, K.M., Biddle, T.M., and Limpus, C.L. (2012) Marine wildlife stranding and mortality database annual report 2008-2010. II. Cetacean and Pinniped. Conservation Technical Data Report, Queensland Government Department of Environment and Heritage Protection.

Melville, A. J. and R. M. Connolly (2003). "Spatial analysis of stable isotope data to determine primary sources of nutrition for fish." *Oecologia* 136: 499-507.

Melville, A. J. and R. M. Connolly (2005). "Food webs supporting fish over subtropical mudflats are based on transported organic matter not in situ microalgae." *Marine Biology* 148: 363-371.

Moriarty, D. (1977). Quantification of carbon, nitrogen and bacterial biomass in the food of some penaeid prawns. *Australian Journal of Marine and Freshwater Research* 28, 113-118.

Morton, R. M., Pollock, B.R., Beumer, J.P., (1987). "The occurrence and diet of fishes in a tidal inlet to a saltmarsh in southern Moreton Bay Qld." *Australian Journal of Ecology* 12: 217-237.

Morton, J. W. (1997). Ecological effects of dredging and dredge spoil disposal. A literature review Technical Paper. US Fish Wildlife Service, 94, 1–33.

Mousalli, A. and R. Connolly (1998). Fish use of the inundated waters of a subtropical saltmarsh-mangrove complex in southeast Queensland. Moreton Bay and Catchment, Brisbane, University of Queensland.

Mustoe, S. (2006). Penguins and marine mammals: final report. Melbourne: AES Applied Ecology Solutions Pty Ltd (through Maunsell Australia Pty Ltd), for the Port of Melbourne Corporation.

Mustoe, S. (2008). Townsville Ocean Terminal: dolphins, dugongs and marine turtles report Ecological impact assessment report. Melbourne: AES Applied Ecological Solutions Pty Ltd, for City Pacific Limited.

NIMPCG. (2011). National System for the Prevention and Management of Marine Pest Incursions. n.s: National Introduced Marine Pests Coordination Group (NIMPCG).

Odum, W. E., C. C. McIvor, T.J. Smith (1985). The ecology of the mangroves of South Florida: a community profile, University of Virginia USA.

Olds, A. (2002). Utilisation by prawns of seagrass beds at different distances from mangrove forests. School of Integrative Biology. Brisbane, University of Queensland.

Olsen, H. F. (1983). Biological Resources of Trinity Inlet and Bay, Queensland.

Olsen, J.L., Stam, W.T., Coyer, J.A., Reusch T.B.H., Billingham, M., Boström C. Calvert, E., Christie, H., Granger, S., Lumière, R. la, Milchakova, N., Oudot-le Secq, M.-P., Procaccini, G., Sanjabi, B., Serrão, E., Veldsink, J., Iddicombe, S., Wyllie-Echeverria, S. (2004). "North Atlantic phylogeography and largescale population differentiation of the seagrass *Zostera marina*." *L. Mol. Ecol.* 13: 1923-1941.

Orth, R. J., Harwell, M. C., & Inglis, G. J. (2006). Ecology of seagrass seeds and seagrass dispersal processes, in Larkum, A. W. D. Orth, R. J. and Duarte, C. M. (eds). Springer, Netherlands. *Seagrasses: Biology, Ecology, and Conservation*.

Parra G.J., Corkeron P.J., and Marsh H. (2004) The Indo-Pacific Humpback Dolphin, *Sousa chinensis* (Osbeck, 1765), in Australian Waters: A Summary of Current Knowledge. *Aquatic Mammals* 2004, 30(1), pp. 197-206. DOI 10.1578/AM.30.1.2004.197

Parra G.J., Corkeron P.J., Marsh H. (2006a) Population sizes, site fidelity and residence patterns of Australian snubfin and Indo-Pacific humpback dolphins: Implications for conservation. *Biological Conservation* 129: 167-180

Parra G.J., Schick R., Corkeron P.J. (2006b) Spatial distribution and environmental correlates of Australian snubfin and Indo-Pacific humpback dolphins. *Ecography* 29: 396-406.

Perry, T.W., (1995). Port of Cairns and Trinity Inlet. Directory of Important Wetlands of Australia. Edited Miller, G.J. and Worland, J.L., 2004.

Poiner, I. R., & Kennedy, R. (1984). Complex patterns of change in the macrobenthos of a large sandbank following dredging. *Community analysis. Marine Biology*, 78: 335-352.

Rasheed, M. (1999) Recovery of experimentally created gaps within a tropical *Zostera capricorni* Aschers.) Seagrass meadow, Queensland Australia. *Journal of Experimental Marine Biology and Ecology*, 235:183-200.

Rasheed, M. (2004). Recovery and succession in a multi-species tropical seagrass meadow following experimental disturbance: the role of sexual and asexual reproduction. *Journal of Experimental Marine Biology and Ecology*, 310, 13-45.

Rasheed, M.A. and Taylor, H.A. (2008). Port of Townsville seagrass baseline survey report, 2007 – 2008. DPI&F Publication PR08-4014 (DPI&F, Cairns), 45 pp.

Rasheed MA, McKenna SA and Tol S (2013). Seagrass habitat of Cairns Harbour and Trinity Inlet: Annual Monitoring and updated Baseline Survey – 2012. JCU Publication, Centre for Tropical Water and Aquatic Ecosystem Research, Cairns.

Robin, J. (1995). Estimated catch and mortality of sea turtles from the East Coast Otter Trawl Fishery of Queensland, Australia. *Biological Conservation*, 74, 157-167.

Skilleter, G. (1998). Ecology of Benthic Invertebrates in Moreton Bay. In: Tibbetts, I.R., Hall, N.J. and Dennison, W.C. (eds) *Moreton Bay and Catchment*. Brisbane: The University of Queensland.

Smith, J. N., Grantham, H., Gales, N., Double, M., Noad, M., & Paton, D. (2012). Identification of humpback whale breeding and calving habitat in the Great Barrier Reef. *Marine Ecology Progress Series* 447, 259-272.

Smith, T. J., and Duke, N.C. (1987). Physical determinants of inter-estuary variation in mangrove species richness around the tropical coastline of Australia. *Journal of Biogeography*, 14(1), 9-19.

Somers, I. (1994). Species composition and distribution of commercial Penaeid prawn catches in the Gulf of Carpentaria, Australia, in relation to depth and sediment type. *Australian Journal of Marine and Freshwater Research*, 45, 317-315.

Souter, D. (2009). Introduced species in the Great Barrier Reef. Reef and Rainforest Research Centre Accessed: July 2011 Available: <http://e-atlas.org.au/content/introduced-species-great-barrier-reef>

Staples, D. J., D. J. Vance, Heals, D.S. (1985). Habitat Requirements of Juvenile Penaeid Prawns and their Relationship to Offshore Fisheries. Second Australian National Prawn Seminar, Kooralbyn, NPS2.

Stevens, J.D., Pillans, R.D. and Salini, J. (2005) Conservation Assessment of *Glyphis sp. A* (spartooth shark), *Glyphis sp. C* (northern river shark), *Pristis microdon* (freshwater sawfish) and *Pristis zijsron* (green sawfish). CSIRO Marine Research, Hobart, Australia.

Swimmer, Y., Arauz, R., Higgins, B., McNaughton, L., McCracken, M., Ballesterio, J., et al. (2005). Food color and marine turtle feeding behavior: Can blue bait reduce turtle bycatch in commercial fisheries? *Marine Ecology Progress Series*, 295, 273-278.

Taylor, H., & Rasheed, M. (2009). Port of Townsville Long-Term Seagrass Monitoring – 2009. Not Stated: Commonwealth of Australia (Department of Employment, Economic Development and Innovation).

Taylor, S., Webley, J., McInnes K. (2012) 2010 Statewide Recreational Fishing Survey. Queensland Government Department of Agriculture, Fisheries and Forestry.

The Coordinator-General (2012) Cairns Shipping Development Project – Terms of reference for an environmental impact statement, November 2012. Prepared by the Queensland Government Office of The Coordinator-General, Brisbane.

UNESCO (United Nations Educational, Scientific and Cultural Organisation) (undated) World heritage Convention: Great Barrier Reef. Accessed: March 2011 Available: [whc.unesco.org/en/list/154](http://whc.unesco.org/en/list/154).

Unsworth R.K.F., Taylor H.A., Rasheed M.A. (2009) Port of Townsville long-term seagrass monitoring: October 2008 QPI&F Publication PR09-4330 (QPI&F, Cairns), 30 pp.

Unsworth, R. K. F., S. A. McKenna, et al. (2010). Seasonal dynamics, productivity and resilience of seagrass at the Port of Abbot Point: 2008 - 2010. Cairns, Fisheries Queensland, DEEDI 68 pp.

Wassenberg, T., & Hill, B. (1987). Natural diet of the Tiger Prawns *Penaeus esculentus* and *P. semisulcatus*. *Australian Journal of Marine and Freshwater Research*, 38, 169-182.

Waycott, M., Longstaff, B.J., Mellors, J. (2005) Seagrass population dynamics and water quality in the Great Barrier Reef region: A review and future research directions. *Marine Pollution Bulletin*, 51: 343–350.

WBM (1997) State of the Environment Report – Trinity Inlet. Prepared for Trinity Inlet Management Program by WBM Oceanics Australia, Brisbane.

WBM. (2004). Moreton Bay Sand Extraction Study Phase 2 - Benthic Fauna Assessments. n.s: WMB, for Moreton Bay Sand Extraction Steering Committee.

Williams, M. (1982). Natural food and feeding in the commercial sand crab *Portunus pelagicus*. *Journal of Experimental Marine Biology and Ecology*. 59, 165-176.

Williams, L. (1997). Queensland's Fisheries Resources Current Condition and Recent Trends 1998 – 1995. n.s: Queensland Department of Primary Industries Information Series QI197007. 101 pp.

Wilson S. and Swan G. (2003) *A Complete Guide to Reptiles of Australia*. Reed New Holland.

Wilson S. (2005) *A Field Guide to Reptiles of Queensland*. (Reprinted 2009) Reed New Holland.

Worley Parsons (2009) Cairns Port Ocean Disposal Site: Benthic Macro-Invertebrate Infauna and Introduced Marine Pest Monitoring Survey 2009. Report prepared for Far North Queensland Ports Corporation by Worley parsons Services Pty Ltd. August 2009.